

# Infinite-Dimensional Game Optimization via Variational Transport

**Lewis Liu**

ALLIS.ALGO2@GMAIL.COM

*Mila & DIRO, University of Montréal, Canada*

**Yufeng Zhang**

YUFENGZHANG2023@U.NORTHWESTERN.EDU

*Northwestern University, United States*

**Zhuoran Yang**

ZRYANG1993@GMAIL.COM

*Princeton University, United States*

**Reza Babanezhad**

BABANEZHAD@GMAIL.COM

*Samsung - SAIT AI Lab Montréal, Canada*

**Zhaoran Wang**

ZHAORANWANG@GMAIL.COM

*Northwestern University, United States*

## Abstract

Game optimization has been extensively studied when decision variables lie in a finite-dimensional space, of which solutions correspond to pure strategies at the Nash equilibrium (NE) and the gradient descent-ascent (GDA) method works widely in practice. In this paper, we consider infinite-dimensional games by a zero-sum distributional optimization problem over a space of probability measures defined on a continuous variable set, which is inspired by finding a mixed NE for finite-dimensional games. We then aim to answer a natural question: *Will GDA-type algorithms still be provably efficient when extended to infinite-dimensional games?* To answer this question, we propose a particle-based variational transport algorithm based on GDA in the functional space. Specifically, the algorithm performs multi-step functional gradient descent-ascent in the Wasserstein space via alternately pushing two sets of particles in the variable space. By characterizing the gradient estimation error from variational form maximization and the convergence behavior of each player with different objective landscapes, we prove rigorously that the generalized GDA algorithm converges to the NE or the value of the game efficiently for a class of games under the Polyak-Łojasiewicz (PL) condition. To the best of our knowledge, we provide the first statistical and convergence guarantees for solving an infinite-dimensional zero-sum game via a provably efficient particle-based method under mild conditions. Additionally, our work provides the first thorough statistical analysis for the particle-based algorithm to learn an objective functional with a variational form using universal approximators (*i.e.*, neural networks (NNs)), which may be of independent interest.

## 1. Introduction

Recent years have witnessed a resurgence in zero-sum games for machine learning applications, where two players' strategies are usually parameterized with two finite-dimensional decision variables. Furthermore, the optimal strategies define the *pure* NE in the sense that they identify two deterministic strategies. Motivating examples include generative adversarial networks (GANs) [23, 47, 57, 59], reinforcement learning [13, 29], distributionally robust optimization (DRO) [22, 62], and learning exponential families [14], among others. Such zero-sum games have been extensively analyzed in convex-concave settings, where a global Nash equilibrium (NE) can be computed efficiently by gradient descent-ascent (GDA) type algorithms [19, 28, 43, 44]. Nonetheless, these methods mostly stagger in nonconvex landscapes for modern applications, for which a bunch of modified gradient-based methods [32, 57, 58] under different settings relax nonconvexity to find a first-order NE [50]. Despite the popularity in exploring variants of the pure NE in nonconvex-nonconcave game settings, another crucial problem arises from a game-theoretic perspective: *What if the pure NE does*

not exist [2, 32]? The finite-dimensional formulation naturally excludes a potential better or even the only existential optimal mixed strategies, and meanwhile is restricted to local convergence without convexity.

To alleviate the concern above and to further understand the difficulty at the boundary of contemporary game optimization, we consider a class of zero-sum infinite-dimensional games where each decision variable is a probability measure representing the mixed strategies over the spaces of pure strategies. In addition, we assume this distributional games to satisfy Riemannian Polyak-Łojasiewicz (PL) conditions and smoothness assumptions, which are satisfied by a range of nonconvex landscapes and the practical training objectives for GANs with regularization [2]. A natural approach to distributional optimization problems is the particle-based method [53, 65, 68], where stochastic gradient Langevin dynamics (SGLD) is usually adopted to draw a sample from the desired distribution via discretization of stochastic differential equations [31]. However, SGLD sampling is quite inefficient for reaching a stationary distribution at each step. Meanwhile, from the view of games, GDA-type algorithms have not been studied in full generality for infinite-dimensional settings. Motivated by the two facts above, we adapt the multi-step GDA-type algorithm to infinite-dimensional games through particle-based approximation, and provide the first set of theoretical guarantees by analyzing its new behavior under infinite-dimensional settings.

We conclude our contributions as follows. (1) To model the mixed NE of finite-dimensional games, we introduce the generic infinite-dimensional zero-sum games. We establish the GDA-type algorithm in the Wasserstein space, also named variational transport for infinite-dimensional games (VTIG), for such games via Riemannian gradient propositions (Proposition 5 and 6). (2) We provide the first thorough analysis of both statistical and optimization errors for VTIG in two scenarios. One is the convergence to the first-order NE under a Riemannian PL condition (Theorem 8), and the other is the convergence to the minimax value under a stronger two-sided PL condition (Theorem 10). (3) As a technical component, we provide statistical analysis for particle-based gradient estimation via an upper bound of the gradient norm by the generalization error of an NN.

**Related work.** Finite-dimensional games under convex-concave settings [28, 33, 43, 44] are adequately studied with corresponding monotonic variational inequalities [15, 23] and solved by GDA [61]. Meanwhile, primal-dual schemes and negative momentum [8, 17, 24] are proposed to help GDA on convergence, which bypasses cyclic dynamics [16, 40, 41]. To tame nonconvexity, [32] proves the  $\mathcal{O}(\theta^{-4})$  rate in gradient evaluations is required in the convergence to an  $\theta$ -first order NE with Max-oracle; [38, 39] reached the same rate when the objective is concave *w.r.t.* the max-player strategy; improved rates of  $\mathcal{O}(\theta^{-3.5})$  and  $\mathcal{O}(\theta^{-2})$  are shown in [58] under PL-game conditions, which is similar to our setting. However, our results are derived for infinite dimensions as a mixed-strategy extension.

In machine learning literature, the notion of mixed NE for GANs was originally presented in [26] without an algorithm to find it. A line of work [2, 27, 31, 48] seeks to further understand and find mixed NEs of GANs. Nonetheless, the existing algorithm in [31] using SGLD is computationally demanding at each step and complicated in the idea of algorithm design without statistical analysis. Our analysis extends the GDA-type algorithm to the Wasserstein space, and shows the existence of a provably efficient particle-based algorithm that pushes a fix-sized set of particles instead of running SGLD repeatedly.

Optimizing functionals of probability measures was studied by Frank-Wolfe [21] and steepest descent algorithms [42] in earlier times. More recently, descent methods in the space of probability measures [20, 54] are getting popular in machine learning, where particle-based methods [9, 36] approximate measures for practical implementation. Similarly, two sets of particles in our algorithm also provide the Dirac measure approximation for probability measures.

**Notations.** We denote by  $[n]$  the set of integers  $\{0, 1, \dots, n\}$ . Let  $\mathcal{C}(\mathbb{R}^d)$  be the set of continuous functions over the  $d$ -dimensional real space  $\mathbb{R}^d$ . Let  $\mathcal{X}$  be a convex compact set in  $\mathbb{R}^d$ . Given a nonnegative measure

$\mu$  on  $\mathcal{X}$ , we define the  $\ell_p$ -norm of the function  $f \in \mathcal{C}(\mathbb{R}^d)$  on  $\mathcal{X}$  as  $\|f\|_{L^p_\mu(\mathcal{X})} = (\int_{\mathcal{X}} |f|^p d\mu)^{1/p}$ . Let  $\mathcal{P}(\mathcal{X})$  denote the collection of all Borel probability measures on the measurable space  $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$ , where  $\mathcal{B}(\mathcal{X})$  is the Borel  $\sigma$ -algebra on  $\mathcal{X}$ . We denote by  $\mathcal{P}_2(\mathcal{X}) \subseteq \mathcal{P}(\mathcal{X})$  the set of Borel probability measures with finite second moments. We define the metric space  $(M, \|\cdot\|)$  by a vector space  $M$  and a metric induced by the norm  $\|\cdot\|$ .

## 2. Problem Formulation and Optimization over Wasserstein Spaces

Below we state the formulation and assumptions for infinite-dimensional games in the Wasserstein space.

### 2.1. From Finite-Dimensional to Infinite-Dimensional Games

Consider the classical formulation of a two-player zero-sum game as follows,

$$\min_{x^\mu \in \mathcal{X}_\mu} \max_{x^\nu \in \mathcal{X}_\nu} f(x^\mu, x^\nu), \quad (1)$$

where  $\mathcal{X}_\mu, \mathcal{X}_\nu \subseteq \mathbb{R}^d$  with  $d > 0$  are convex compact sets of pure strategies as  $d$ -dimensional vectors with periodic or zero-flux boundary conditions *w.r.t.* the vector fields specified in Proposition 6, and  $f$  is the objective function. In nonconvex-nonconcave regimes, as finding local Nash equilibria is NP-hard or even impossible [32], a weaker notion of *first-order* NE (FNE) for a pair  $(x_*^\mu, x_*^\nu) \in \mathcal{X}_\mu \times \mathcal{X}_\nu$  is defined as

$$\begin{aligned} \langle \nabla_{x^\mu} f(x_*^\mu, x_*^\nu), x^\mu - x_*^\mu \rangle &\geq 0, \\ \langle \nabla_{x^\nu} f(x_*^\mu, x_*^\nu), x^\nu - x_*^\nu \rangle &\leq 0, \quad \forall x^\mu \in \mathcal{X}_\mu, \forall x^\nu \in \mathcal{X}_\nu, \end{aligned} \quad (2)$$

which corresponds to first-order necessary optimality conditions. Observing that without a probability representation (1) only admits pure Nash strategies, we lift (1) by considering distributions over  $\mathcal{X}_\mu$  and  $\mathcal{X}_\nu$  to allow mixed strategies. The infinite-dimensional distributional two-player zero-sum game is defined as

$$\min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} \max_{\nu \in \mathcal{M}(\mathcal{X}_\nu)} F(\mu, \nu). \quad (3)$$

Here  $F: \mathcal{M}(\mathcal{X}_\mu) \times \mathcal{M}(\mathcal{X}_\nu) \rightarrow \mathbb{R}$  is the objective functional. Also,  $\mathcal{M}(\mathcal{X}) = (\mathcal{P}_2(\mathcal{X}), \mathcal{W}_2)$  is the Wasserstein ( $W_2$ -) space, an infinite-dimensional manifold by [64], with the  $W_2$ -distance on  $\mathcal{P}_2(\mathcal{X})$  defined as

$$\mathcal{W}_2(\mu, \nu) = \inf \left\{ \mathbb{E}[\|X - Y\|^2]^{1/2} \mid \mathcal{L}(X) = \mu, \mathcal{L}(Y) = \nu \right\},$$

where the infimum is taken over the random variables  $X$  and  $Y$ , and we denote by  $\mathcal{L}(X)$  the law of a random variable  $X$ . Without loss of generality, we set  $\mathcal{X}_\mu = \mathcal{X}_\nu = \mathcal{X}$  and write  $\mathcal{M} = \mathcal{M}(\mathcal{X})$ . Without specification, the domain of an integral is the set  $\mathcal{X}$ . We refer to the two players as player  $\mu$  and player  $\nu$ , respectively. To further characterize the properties of such a Wasserstein space  $\mathcal{M}$ , we introduce geodesics, tangent vectors and tangent spaces below.

**Definition 1** *Define the smooth curve  $\gamma: [0, 1] \rightarrow \mathcal{P}_2(\mathcal{X})$ . We call the curve  $\gamma$  a geodesic if there exists a constant  $v \geq 0$  such that  $\|\gamma(t_1) - \gamma(t_2)\| = v \cdot |t_1 - t_2|$  for any  $t_1, t_2 \in [0, 1]$ . A tangent vector at  $\mu \in \mathcal{M}$  is an equivalence class of differentiable curves through  $\mu$  with a prescribed velocity vector at  $\mu$ . The tangent space at  $\mu$ , denoted by  $T_\mu \mathcal{M}$ , consists of all tangent vectors at  $\mu$ .*

Furthermore, the manifold  $\mathcal{M}$  is equipped with a weak Riemannian structure in the following sense [64]. Given any tangent vectors  $u, v$  at  $\mu \in \mathcal{M}$  and the vector fields  $\tilde{u}, \tilde{v}$  satisfying continuity equations  $u = -\text{div}(\mu \tilde{u})$  and  $v = -\text{div}(\mu \tilde{v})$ , respectively, we define the inner product of  $u$  and  $v$  as  $\langle u, v \rangle_\mu = \int \langle \tilde{u}, \tilde{v} \rangle d\mu$ , where  $\langle \tilde{u}, \tilde{v} \rangle$  is the inner product in  $\mathcal{X}$ . Such a metric induces a norm  $\|u\|_\mu = \langle u, u \rangle_\mu^{1/2}$  for any  $u \in T_\mu \mathcal{M}$ .

Under such a structure, we define the directional derivative *w.r.t.*  $u \in T_\mu \mathcal{M}$  of a differentiable functional  $g: \mathcal{M} \rightarrow \mathbb{R}$  as  $\nabla_u g(\mu) = \left. \frac{d}{dt} g[\gamma(t)] \right|_{t=0}$ , where  $\gamma(0) = \mu \in \mathcal{M}$ ,  $\gamma'(0) = u$ . In addition, we say  $g$  is  $W_2$ -differentiable at  $\mu$  if there exists  $u' \in T_\mu \mathcal{M}$  such that  $\nabla_u g(\mu) = \langle u', u \rangle_\mu$  for any  $u \in T_\mu \mathcal{M}$ , and write  $\text{grad } g(\mu) = u'$  as the (weak) Riemannian gradient of  $g$  at  $\mu$ . The partial gradient  $\text{grad}_\mu F(\mu, \nu)$  is defined similarly for a functional  $F: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$  when fixing  $\nu$ . The exponential map at  $\mu$ , denoted by  $\text{Exp}_\mu$ , sends any  $u \in T_\mu \mathcal{M}$  to  $\mu' = \gamma_u(1)$ <sup>1</sup>, where  $\gamma_u$  is a geodesic such that  $\gamma_u(0) = \mu$ ,  $\gamma'_u(0) = u$ . For any  $\mu, \nu \in \mathcal{M}$ , the parallel transport  $\Gamma_\mu^\nu: T_\mu \mathcal{M} \rightarrow T_\nu \mathcal{M}$  is the map such that  $\langle u, v \rangle_\mu = \langle \Gamma_\mu^\nu u, \Gamma_\mu^\nu v \rangle_\nu$  for any  $u, v \in T_\mu \mathcal{M}$ . Also, as  $\mathcal{X}$  is separable and complete,  $\mathcal{M}$  is geodesically complete [63] in the sense that the exponential map is defined on the whole tangent space. See §B for more formal definitions.

We also assume the objective functional  $F$  in (3) to satisfy the following variational forms,

$$F(\mu, \nu) = F_\nu(\mu) = \sup_{f \in \mathcal{F}} \left\{ \int_{\mathcal{X}} f d\mu - F_\nu^*(f) \right\}, \quad F(\mu, \nu) = F_\mu(\nu) = - \sup_{f \in \mathcal{F}} \left\{ \int_{\mathcal{X}} f d\nu - F_\mu^*(f) \right\}, \quad (4)$$

where  $\mathcal{F}$  is the family of square-integrable functions over  $\mathcal{X}$ ,  $F_\mu^*: \mathcal{F} \rightarrow \mathbb{R}$  and  $F_\nu^*: \mathcal{F} \rightarrow \mathbb{R}$  are strongly convex and smooth functional *w.r.t.* the  $\ell_2$ -norm. In fact, (4) follows from definitions of the conjugate function, and the example in §2 shows that a wide class of  $f$ -divergences admits such forms.

For theoretical analysis, we impose the following assumptions on the objective functional  $F$ .

**Assumption 2** *We assume that  $F$  is Lipschitz continuous and smooth w.r.t. the Wasserstein distance in the sense that*

$$\begin{aligned} |F(\mu_1, \nu) - F(\mu_2, \nu)| &\leq L_\mu \mathcal{W}_2(\mu_1, \mu_2), \\ |F(\mu, \nu_1) - F(\mu, \nu_2)| &\leq L_\nu \mathcal{W}_2(\nu_1, \nu_2), \\ \mathbf{d}(\text{grad } F_\nu(\mu_1), \text{grad } F_\nu(\mu_2)) &\leq L_1 \cdot \mathcal{W}_2(\mu_1, \mu_2), \quad \mathbf{d}(\text{grad } F_\mu(\nu_1), \text{grad } F_\mu(\nu_2)) \leq L_2 \cdot \mathcal{W}_2(\nu_1, \nu_2), \\ \mathbf{d}(\text{grad } F_{\mu_1}(\nu), \text{grad } F_{\mu_2}(\nu)) &\leq L_0 \cdot \mathcal{W}_2(\mu_1, \mu_2), \quad \mathbf{d}(\text{grad } F_{\nu_1}(\mu), \text{grad } F_{\nu_2}(\mu)) \leq L_0 \cdot \mathcal{W}_2(\nu_1, \nu_2) \end{aligned} \quad (5)$$

for any  $\mu, \mu_1, \mu_2, \nu, \nu_1, \nu_2 \in \mathcal{M}$ . Here  $L_\mu, L_\nu, L_1, L_2$ , and  $L_0$  are absolute constants and  $\mathbf{d}^2(u, v) = \langle u - \Gamma_\nu^\mu v, u - \Gamma_\nu^\mu v \rangle_\mu$  for any  $\mu, \nu \in \mathcal{M}$ ,  $u \in T_\mu \mathcal{M}$ , and  $v \in T_\nu \mathcal{M}$ .

Assumption 2 is a natural extension of Lipschitz continuity and smoothness for Euclidean space to  $\mathcal{M}$ , where the Euclidean distance is replaced by  $\mathcal{W}_2$ . The following assumption extends the notion of PL condition, also known as gradient domination [45, 52, 58], to infinite-dimensional spaces.

**Assumption 3** (Riemannian PL condition). *A  $W_2$ -differentiable functional  $g: \mathcal{M} \rightarrow \mathbb{R}$  with minimum value  $g^* = \inf_{\mu \in \mathcal{M}} g(\mu)$  is called  $\xi$ -PL ( $\xi$ -gradient dominated) if for all  $\mu \in \mathcal{M}$  we have*

$$\langle \text{grad } g(\mu), \text{grad } g(\mu) \rangle_\mu \geq 2\xi (g(\mu) - g^*). \quad (6)$$

We call (3) a  $\xi$ -PL game, or simply a PL game, if  $H_\mu(\nu) \triangleq -F(\mu, \nu)$  is  $\xi$ -PL. We assume (3) to be a  $\xi$ -PL game.

In particular, Assumption 3 implies that if the norm of the gradient is small at  $\mu \in \mathcal{M}$ , then the functional value at  $\mu$  will be close to the optimum. In addition, it is not restrictive since a non-convex functional can still satisfy the PL condition [34]. To justify all the above assumptions, we provide the following example stemming from learning GANs, where the pure strategies in (3) correspond to parameters  $x^\mu$  and  $x^\nu$  of the GAN.

1. Hence, for  $\mu_1, \mu_2 \in \mathcal{M}$ ,  $\text{Exp}_{\mu_1}^{-1}(\mu_2)$  is an analogy to  $x_2 - x_1$  for  $x_1, x_2 \in \mathcal{X}$ .

**Example 1** Consider the mixed NE of WGANs [1] with Kullback-Leibler (KL) divergence regularization,

$$\min_{\mu \in \mathcal{M}} \max_{\nu \in \mathcal{M}} \mathbb{E}_{x^\nu \sim \nu} \mathbb{E}_{\zeta \sim \mathbb{P}_{real}} [h_{x^\nu}(\zeta)] - \mathbb{E}_{x^\nu \sim \nu} \mathbb{E}_{x^\mu \sim \mu} \mathbb{E}_{\zeta \sim \mathbb{P}_{x^\mu}} [h_{x^\nu}(\zeta)] - \text{KL}(\nu \parallel \mu_0) + \text{KL}(\mu \parallel \mu_0), \quad (7)$$

where  $\text{KL}(\mu \parallel \lambda) = \int_{\mathcal{X}} \log(d\mu/d\lambda) d\mu$  with Lebesgue measures  $\mu$  and  $\lambda$ , and  $\mu_0$  is the probability measure of standard Gaussian. Also,  $h_\nu$  denotes the discriminator parameterized by NNs, of which the input is  $\zeta \in \mathcal{X}$ . Without the expectations of  $x^\mu$  and  $x^\nu$ , (7) is reduced to the original regularized WGAN objective that admits only finite-dimensional pure Nash strategies. Further, we define the linear operator  $D : \mathcal{M} \rightarrow \mathcal{F}$  by  $(D\mu)(x^\nu) = \mathbb{E}_{x^\mu \sim \mu} \mathbb{E}_{\zeta \sim \mathbb{P}_{x^\mu}} [h_{x^\nu}(\zeta)]$  for any  $x^\nu \in \mathcal{X}$  and some continuous function  $h_{x^\nu} \in \mathcal{F}$ . We also define  $g(x^\nu) = \mathbb{E}_{\zeta \sim \mathbb{P}_{real}} [h_{x^\nu}(\zeta)]$ . Then the objective  $F$  in (7) can be rewritten as  $F(\mu, \nu) = \langle \nu, g \rangle - \langle \nu, D\mu \rangle - \text{KL}(\nu \parallel \mu_0) + \text{KL}(\mu \parallel \mu_0)$ . It follows from the logarithmic Sobolev inequality (LSI) [49] in  $W_2$ -space that player  $\mu$  meets the PL condition. Since the KL divergence is an  $f$ -divergence, the variational forms are guaranteed as follows,

$$F_\nu(\mu) = \sup_{f \in \mathcal{F}} \left\{ \int f d\mu - \int \exp \{ f(x^\mu) + \mathbb{E}_{x^\nu \sim \nu} \mathbb{E}_{\zeta \sim \mathbb{P}_{x^\mu}} [h_{x^\nu}(\zeta)] \} d\mu_0(x^\mu) + \widehat{F}_\nu \right\}, \quad (8)$$

$$F_\mu(\nu) = - \sup_{f \in \mathcal{F}} \left\{ \int f d\nu - \int \exp \{ f(x^\nu) + g(x^\nu) - (D\mu)(x^\nu) \} d\mu_0(x^\nu) + \widehat{F}_\mu \right\}. \quad (9)$$

Here  $\widehat{F}_\nu = 1 - \text{KL}(\nu \parallel \mu_0) + \mathbb{E}_{x^\nu \sim \nu} \mathbb{E}_{\zeta \sim \mathbb{P}_{real}} [h_{x^\nu}(\zeta)]$  and  $\widehat{F}_\mu = 1 - \text{KL}(\mu \parallel \mu_0)$  are constants when fixing  $\nu$  and  $\mu$ , respectively. See §C.3 for details. We remark that in practical GAN training, KL regularization terms exist to prevent the mode collapse. More generally, the KL-regularized distributional bilinear game  $\min_{\mu \in \mathcal{M}} \max_{\nu \in \mathcal{M}} \langle \nu, A\mu \rangle - \text{KL}(\nu \parallel \mu_0) + \text{KL}(\mu \parallel \mu_0)$  given a linear operator  $A : \mathcal{M} \rightarrow \mathcal{F}$  is widely studied in games. Similarly, we write its variational forms as  $F_\nu(\mu) = \sup_{f \in \mathcal{F}} \{ \int f d\mu - \int \exp \{ f(x^\nu) - A^* \nu(x^\nu) \} d\mu_0(x^\nu) + 1 - \text{KL}(\nu \parallel \mu_0) \}$  and  $F_\mu(\nu) = - \sup_{f \in \mathcal{F}} \{ \int f d\nu - \int \exp \{ f(x^\mu) + A\mu(x^\mu) \} d\mu_0(x^\mu) + 1 - \text{KL}(\mu \parallel \mu_0) \}$ , where  $A^*$  is the adjoint of  $A$ .

## 2.2. Measurement of Solutions

To quantify the accuracy of solutions to (3), we generalize the NE of finite-dimensional games to our infinite-dimensional distributional games. Given the numerical accuracy of iterative algorithms in practice, we define the notion of infinite-dimensional first-order NE (IFNE) as a performance measure.

**Definition 4 (IFNE)** For any  $\mu_1, \nu_1 \in \mathcal{M}$ , we define

$$\begin{aligned} \mathcal{J}_\mu(\mu_1, \nu_1) &\triangleq - \min_{W_2(\mu, \mu_1) \leq 1} \langle \text{grad}_\mu F(\mu_1, \nu_1), \text{Exp}_{\mu_1}^{-1}(\mu) \rangle_{\mu_1}, \\ \mathcal{J}_\nu(\mu_1, \nu_1) &\triangleq \max_{W_2(\nu, \nu_1) \leq 1} \langle \text{grad}_\nu F(\mu_1, \nu_1), \text{Exp}_{\nu_1}^{-1}(\nu) \rangle_{\nu_1} \end{aligned}$$

as the first-order errors (FEs). Then a point  $(\mu^*, \nu^*) \in \mathcal{M} \times \mathcal{M}$  is called a  $\theta$ -IFNE of (3) if

$$\mathcal{J}_\mu(\mu^*, \nu^*) \leq \theta \quad \text{and} \quad \mathcal{J}_\nu(\mu^*, \nu^*) \leq \theta. \quad (10)$$

When  $\theta = 0$ , we call  $(\mu^*, \nu^*)$  an IFNE. Definition 4 characterizes how far the solutions are from the FNE in the  $W_2$ -space. Also, we characterize the upper bound  $\theta$  in terms of the problem parameters for convergence rates in §4.

### 3. Variational Transport Algorithm for Infinite-Dimensional Games

In what follows, we introduce the variational transport algorithm to characterize GDA for the infinite-dimensional game defined in (3). Our idea is based on the multi-step GDA algorithm in [58] with nested loops, where multiple gradient ascent steps are run for estimating the gradient of the *inner maximization functional* defined as  $G(\mu) = \max_{\nu \in \mathcal{M}} F(\mu, \nu)$  w.r.t.  $\mu$ , which provides a descent direction for the outer minimization problem. Without specifying, statements below hold for both  $\mu$  and  $\nu$  although they are presented by  $\mu \in \mathcal{M}$ .

#### 3.1. Gradient Descent beyond the Euclidean Space

We first show the connection between functional gradient descent in the Wasserstein Space  $\mathcal{M}$  and transportation maps in the variable space  $\mathcal{X}$ . Specifically, we expect to update the current iterate  $\mu \in \mathcal{M}$  of the gradient descent in the direction of  $\text{grad } F_\nu(\mu)$  along the geodesic. Therefore, in the ideal case, the gradient update is given by

$$\mu \leftarrow \text{Exp}_\mu[-\eta \cdot \text{grad } F_\nu(\mu)], \quad (11)$$

where  $\eta > 0$  is the stepsize. The proposition below bridges the Riemannian gradient of a  $W_2$ -differentiable functional on  $\mathcal{M}$  and its functional gradient w.r.t. the  $\ell_2$ -norm. We denote by  $f_\mu^* \in \mathcal{F}$  the optimal solution to (4) for  $F_\nu(\mu)$ .

**Proposition 5 (Riemannian Gradients to Functional Gradients)** *Let  $F : \mathcal{M} \rightarrow \mathbb{R}$  be a  $W_2$ -differentiable functional, with its functional gradient w.r.t. the  $\ell_2$ -norm written as  $\delta F / \delta \mu$ . Then, it follows that*

$$\text{grad } F(\mu) = -\text{div} \left[ \mu \cdot \nabla \left( \frac{\delta F}{\delta \mu} \right) \right], \quad (12)$$

where  $\text{div}$  is the divergence operator on  $\mathcal{X}$ . Furthermore, by the variational form of (3), we have  $\delta F_\nu / \delta \mu = f_\mu^*$  and  $\text{grad } F_\nu(\mu) = -\text{div}(\mu \cdot \nabla f_\mu^*)$ .

**Proof** See §C.1 for a detailed proof. ■

By Proposition 5, to obtain a descent direction in  $W_2$ -space for  $F_\nu(\mu)$ , we first solve (4) for  $f_\mu^* \in \mathcal{F}$  and then, compute the divergence in (12). Also,  $\text{Exp}_\mu$  in (11) needs to be specified. As in practice we only have access to samples, or particles, from  $\mu$ , we establish the proposition below to perform approximate gradient updates in (11) via particles.

**Proposition 6 (Pushing particles as an exponential map)** *For any  $\mu \in \mathcal{M}$  and any  $s \in T_\mu \mathcal{M}$ , suppose the elliptic equation  $-\text{div}(\mu \cdot \nabla u) = s$  admits a unique solution  $u : \mathcal{X} \rightarrow \mathbb{R}$  such that  $\nabla u : \mathcal{X} \rightarrow \mathbb{R}^d$  is  $h$ -Lipschitz continuous. Then, for any  $t \in [0, 1/h)$ , we have*

$$[\text{Exp}_\mathcal{X}(t \cdot \nabla u)]_\# \mu = \text{Exp}_\mu(t \cdot s), \quad (13)$$

where we use  $\text{Exp}_\mathcal{X}(t \cdot \nabla u)$  to denote the transportation map on  $\mathcal{X}$  that sends any  $x \in \mathcal{X}$  to a point  $\text{Exp}_x(t \cdot \nabla u(x)) \in \mathcal{X}$ , which is also the exponential map over  $\mathcal{X}$ . We denote by  $T_\# : \mathcal{P}_2(\mathcal{X}) \rightarrow \mathcal{P}_2(\mathcal{X})$  the push-forward map of a transportation map  $T : \mathcal{X} \rightarrow \mathcal{X}$  such that for any  $\mu \in \mathcal{M}$  and any measurable set  $A \in \mathcal{X}$ , we have  $T_\# \mu(A) = \mu(T^{-1}(A))$ .

**Proof** See §C.2 for a detailed proof. ■

Hence, if  $\nabla f_\mu^*$  is  $h$ -Lipschitz, by Proposition 5 and 6, for any  $t \in [0, 1/h)$  we obtain  $\text{Exp}_\mu[-t \cdot \text{grad } F(\mu)] = [\text{Exp}_\mathcal{X}(-t \cdot \nabla f_\mu^*)]_\# \mu$ . given  $\mu \in \mathcal{M}$ . This identifies the gradient descent update in the Wasserstein spaces




Figure 1: Equivalence between particle pushing in the Euclidean space  $\mathcal{X}$  and the exponential map in the Wasserstein space  $\mathcal{M}$ . The tangent vector  $v \in T_x \mathcal{M}$  at  $x$  induces the exponential map  $\text{Exp}_x$  and its correspondence in  $\mathcal{X}$ , the push-forward map  $T_{\#} = [\text{Exp}_{\mathcal{X}}(-t \cdot \nabla f_{\mu}^*)]_{\#}$ .  $X_k^{\mu}$  is the set of  $\mu$ -particles at timestep  $k$  in Algorithm 1.

with the push-forward map of probability measures over the Euclidean space, which can be approximated by pushing a set of particles. We illustrate such correspondence in Figure 1.

Further, we are left with the variational form maximization (VFM) problem in (4), where difficulties lie in the following aspects. (i) Firstly, our approach is expected to provide the reasonable statistical error incurred by estimating  $f_{\mu}^*$  by  $\tilde{f}_{\mu}^*$  from the empirical version of VFM,

$$\tilde{f}_{\mu}^* = \operatorname{argmax}_{f \in \mathcal{F}} \left\{ \int_{\mathcal{X}} f \, d\hat{\mu} - F_{\nu}^*(f) \right\} = \operatorname{argmax}_{f \in \mathcal{F}} \left\{ \frac{1}{N} \sum_{i=1}^N f(x_i) - F_{\nu}^*(f) \right\}, \quad (14)$$

where we replace  $\mu$  in (4) by the empirical measure  $\hat{\mu} = 1/N \sum_{i=1}^N \delta_{x_i}$ , *i.e.*, an average of Dirac measures over samples  $x_i$ 's. (ii) Secondly, maximization over  $\mathcal{F}$  is computationally intractable. To this end, we perform stochastic gradient descent (SGD) to learn  $f_{\mu}^*$  from the following class  $\tilde{\mathcal{F}}$  of neural networks (NNs) instead of  $\mathcal{F}$ , which is a rich class by the universal approximation theorem [12, 30].

**Neural Network Parametrization.** We consider the following class of NNs,

$$\tilde{\mathcal{F}} = \left\{ \tilde{f} \mid \tilde{f}(x) = \frac{1}{\sqrt{w}} \sum_{i=1}^w b_i \cdot \sigma([\beta]_i^{\top} x) \right\}, \quad (15)$$

where  $w$  is the width of the neural network,  $[\beta]_i \in \mathbb{R}^d$ ,  $\beta = ([\beta]_1^{\top}, \dots, [\beta]_w^{\top})^{\top} \in \mathbb{R}^{wd}$  are input weights,  $\sigma(\cdot)$  denotes a smooth activation function, and  $b_i \in \{-1, 1\}$  ( $i \in [w]$ ) are the output weights. As shown in Algorithm 3, only  $\beta$  is updated during training while  $b_i$  ( $i \in [w]$ ) is fixed. In addition, at each iteration we project the input weights  $\beta$  to an  $\ell_2$ -ball centered at  $\beta(0)$  with radius  $r_f$  defined as  $\mathcal{B}^0(r_f) = \{\beta : \|\beta - \beta(0)\|_2 \leq r_f\}$ . See §D.1 for more details of  $\tilde{\mathcal{F}}$ .

### 3.2. Algorithm for Two-Player Infinite-Dimensional Games

We now put together two nested loops of gradient descent/ascent updates approximated by particles as the variational transport algorithm for infinite-dimensional games (VTIG) in Algorithm 1. In detail, we maintain two sets of  $N_{\mu}$   $\mu$ -particles and  $N_{\nu}$   $\nu$ -particles for player  $\mu$  and player  $\nu$ . Also, VTIG output the

corresponding probability measures approximated by these two sets as the solutions to (3), respectively. At outer-loop timestep  $k$ , we denote the set for player  $\mu$  by  $X_k^\mu = \{x_{i,k}^\mu\}_{i \in [N_\mu]}$  and the set for player  $\nu$  at inner-loop timestep  $l$  of outer-loop timestep  $k$  by  $X_l^\nu(\tilde{\mu}_k) = \{x_{i,l}^\nu(\tilde{\mu}_k)\}_{i \in [N_\nu]}$ . Here we write  $X_l^\nu(\tilde{\mu}_k)$  and  $x_{i,l}^\nu(\tilde{\mu}_k)$  to emphasize that we fix  $X_k^\mu$  (resp.  $\tilde{\mu}_k$ ) when updating  $X_l^\nu$  (resp.  $\tilde{\nu}_l$ ) in Line 7 of Algorithm 1. Also,  $\{\tilde{\mu}_k\}_{k \geq 0}$  and  $\{\tilde{\nu}_l(\tilde{\mu}_k)\}_{k,l \geq 0}$  are sequences of probability measures of  $\{X_k^\mu\}_{k \geq 0}$  and  $\{X_l^\nu(\tilde{\mu}_k)\}_{k,l \geq 0}$  constructed implicitly by VTIG, which is specified later. Further, the set of  $\mu$ -particles  $X_k^\mu$  is updated as follows given  $X_0^\mu$  for  $k \geq 1$ . At the outer-loop timestep  $k$ , VTIG computes the solution to (14) following Line 10 in Algorithm 1

$$\tilde{f}_k^* \leftarrow \mathbf{VFM}(X_k^\mu, F_{\tilde{\nu}_{k+1}}^*, N_\mu) \quad (16)$$

based on the current  $\mu$ -particle set  $X_k^\mu$ , the functional  $F_{\tilde{\nu}_{k+1}}^*$  defined in the variational form (4), and the number of  $\mu$ -particles  $N_\mu$ . As shown in Algorithm 3, the VFM problem is solved by learning a neural network  $\tilde{f}_k^*$  belonging to the class  $\tilde{\mathcal{F}}$  defined in (15) via SGD. With the obtained  $\nabla \tilde{f}_k^*$ , VTIG push  $\mu$ -particles in this direction as follows (Line 11 of Algorithm 1),

$$x_{i,k+1}^\mu \leftarrow \text{Exp}_{x_{i,k}^\mu} \left[ -\eta^\mu \cdot \nabla \tilde{f}_k^*(x_{i,k}^\mu) \right] \quad (17)$$

for all  $i \in [N_\mu]$ . Here  $\eta^\mu > 0$  are the stepsize specified in Theorem 8. This is equivalent to updating the empirical measure  $\hat{\mu} = N_\mu^{-1} \sum_{i \in [N_\mu]} \delta_{x_{i,k}}$  by the pushforward measure  $[\text{Exp}_{\mathcal{X}}(-\eta^\mu \cdot \nabla \tilde{f}_{l,k}^*)]_{\#} \hat{\mu}$ , which approximates the Riemannian gradient update in (11) with stepsize  $\eta^\mu$ . Also, the exponential map in Euclidean space is reduced to a gradient descent step on  $x_{i,k}^\mu \in \mathbb{R}^d$ .

Similarly, to update the set of  $\nu$ -particles  $X_l^\nu(\tilde{\mu}_k)$ , VTIG computes the solution to (14) following Line 6 in Algorithm 1 at inner-loop timestep  $l$  of outer-loop timestep  $k$ . Then, the  $\nu$ -particles are pushed by

$$x_{i,l+1}^\nu(\tilde{\mu}_k) \leftarrow \text{Exp}_{x_{i,l}^\nu(\tilde{\mu}_k)} \left[ \eta^\nu \cdot \nabla \tilde{f}_{l,k}^*(x_{i,l}^\nu(\tilde{\mu}_k)) \right] \quad (18)$$

for all  $i \in [N_\nu]$  in Line 7 of Algorithm 1, with fixed  $\tilde{\mu}_k$ . In particular, the sequences of probability measures  $\{\tilde{\mu}_k\}_{k \geq 0}$  and  $\{\tilde{\nu}_l(\tilde{\mu}_k)\}_{k,l \geq 0}$  are constructed as below. We define sequences of transportation maps  $\{T_k^\mu: \mathcal{X} \rightarrow \mathcal{X}\}_{k=0}^{K_\mu}$  with  $T_0^\mu = \text{id}$  and  $\{T_m^\nu: \mathcal{X} \rightarrow \mathcal{X}\}_{m=0}^{K_\mu K_\nu}$  with  $T_0^\nu = \text{id}$ , by

$$T_{k+1}^\mu = \text{Exp}_{\mathcal{X}}(-\eta^\mu \cdot \nabla \tilde{f}_k^*) \circ T_k^\mu \quad \text{and} \quad T_{kl+1}^\nu = \text{Exp}_{\mathcal{X}}(-\eta^\nu \cdot \nabla \tilde{f}_{l,k}^*) \circ T_{kl}^\nu, \quad (19)$$

respectively for  $k \in [K_\mu], l \in [K_\nu]$ . Here  $K_\mu$  and  $K_\nu$  are the numbers of timesteps of the inner and outer loops, respectively. Then for each  $k \geq 1$  we define  $\tilde{\mu}_k = (T_k^\mu)_{\#} \tilde{\mu}_0$  and  $\tilde{\nu}_k = (T_k^\nu)_{\#} \tilde{\nu}_0$ , where  $\mu_0$  and  $\nu_0$  are initial probability measures. Hence, we have  $\tilde{\nu}_l(\tilde{\mu}_k) = \tilde{\nu}_{lk}$ . Also,  $x_{i,k}^\mu \stackrel{\text{i.i.d.}}{\sim} \tilde{\mu}_k$  and  $x_{i,l}^\nu(\tilde{\mu}_k) \stackrel{\text{i.i.d.}}{\sim} \tilde{\nu}_l(\tilde{\mu}_k)$  are independent samples. Such implicit construction of transportation maps and probability measures also induces a theoretical version of VTIG via resampling. See Algorithm 2 for details. Additionally, we adopt the constructed measure  $\tilde{\nu}_{k+1}$  to compute  $F_{\tilde{\nu}_{k+1}}^*$  in (16) since for most objectives  $F$  such as that in Example 1, we can always sample many enough particles to approximate the expectation terms *w.r.t.*  $\tilde{\nu}_{k+1}$  for  $k \geq 0$ .

#### 4. Main Results

To ensure the independence of the particles for statistical analysis, we adopt Algorithm 2 for theoretical analysis. We characterize the statistical error induced by estimating Riemannian gradients using finite particle samples in §4.1 for both players. In §4.2 we establish the convergence rate of VTIG to the IFNE under the PL condition for one player. Furthermore, we present in §4.3 that under a stronger assumption on the objective  $F$ , *i.e.*, the two-sided PL condition, a linear convergence rate to the minimax value of the game is achieved.



---

**Algorithm 1** Multi-Step Variational Transport Algorithm for Infinite-Dimensional Games (VTIG)

---

- 1: **Input:** Functional  $F: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$ ; initial probability measures  $\tilde{\mu}_0, \tilde{\nu}_0 \in \mathcal{M}$ ; numbers of particles  $N_\mu, N_\nu$ ; numbers of iterations  $K_\mu, K_\nu$ ; and stepsizes  $\eta^\mu \in (0, \min\{1/h, 2/\tilde{L}\})$ ,  $\eta^\nu \in (0, \min\{1/(4L_2), 1/h\})$ .
  - 2: Initialize  $N_\mu$  ( $N_\nu$ ) particles  $X_0^\mu = \{x_{i,0}^\mu\}_{i \in [N_\mu]}$  ( $X_0^\nu$ ) by drawing  $N_\mu$  ( $N_\nu$ ) i.i.d. samples from  $\tilde{\mu}_0$  ( $\tilde{\nu}_0$ ).
  - 3: **for**  $k = 0, 1, 2, \dots, K_\mu - 1$  **do**
  - 4:   Set  $X_0^\nu(\tilde{\mu}_k) = X_{K_\nu}^\nu$
  - 5:   **for**  $l = 0, 1, 2, \dots, K_\nu - 1$  **do**
  - 6:      $\tilde{f}_{l,k}^* \leftarrow \mathbf{VFM}(X_l^\nu(\tilde{\mu}_k), F_{\tilde{\mu}_k}^*, N_\nu)$
  - 7:     Push  $\nu$ -particles:  $x_{i,l+1}^\nu(\tilde{\mu}_k) \leftarrow \text{Exp}_{x_{i,l}^\nu(\tilde{\mu}_k)}[-\eta^\nu \cdot \nabla \tilde{f}_{l,k}^*(x_{i,l}^\nu(\tilde{\mu}_k))]$  for all  $i \in [N_\nu]$
  - 8:   **end for**
  - 9:   Set  $X_{k+1}^\nu = X_{K_\nu}^\nu(\tilde{\mu}_k)$
  - 10:  $\tilde{f}_k^* \leftarrow \mathbf{VFM}(X_k^\mu, F_{\tilde{\nu}_{k+1}}^*, N_\mu)$
  - 11: Push  $\mu$ -particles:  $x_{i,k+1}^\mu \leftarrow \text{Exp}_{x_{i,k}^\mu}[-\eta^\mu \cdot \nabla \tilde{f}_k^*(x_{i,k}^\mu)]$  for all  $i \in [N_\mu]$
  - 12: Set  $X_{k+1}^\mu = \{x_{i,k+1}^\mu\}_{i \in [N_\mu]}$
  - 13: **end for**
  - 14: **Output:**  $\tilde{\mu}^* = N_\mu^{-1} \sum_{i \in [N_\mu]} \delta_{x_{i,K_\mu}}$ ,  $\tilde{\nu}^* = N_\nu^{-1} \sum_{i \in [N_\nu]} \delta_{x_{i,K_\nu}}$
- 

#### 4.1. Statistical Analysis

For each player, VTIG can be viewed as a Riemannian gradient descent method with biased gradient estimates. We characterize the bias in terms of the generalization error of function approximators, where lie the essential difficulties in theory. In this section, we present the analysis for player  $\mu$ . The analysis of player  $\nu$  is similar.

**Gradient estimation.** Recall that by Proposition 5, the desired descent direction for timestep  $k \geq 0$  is  $\text{grad } F(\tilde{\mu}_k) = -\text{div}(\tilde{\mu}_k \cdot \nabla f_k^*)$ . However, with only finite samples, we obtain an estimator  $\tilde{f}_k^*$  of  $f_k^*$ . Hence, the gradient estimate at  $\tilde{\mu}_k$  is  $-\text{div}(\tilde{\mu}_k \cdot \nabla \tilde{f}_k^*)$ , and the difference between  $\text{grad } F(\tilde{\mu}_k)$  and its estimate is denoted by  $\delta_k = -\text{div}[\tilde{\mu}_k \cdot (\nabla \tilde{f}_k^* - \nabla f_k^*)]$ . By observing that  $\delta_k \in T_{\tilde{\mu}_k} \mathcal{M}$  and that the randomness of  $\delta_k$  comes from the initial samples  $X_0^\mu$ , we define

$$\bar{\varepsilon}_k = \mathbb{E}_{X_0^\mu} \langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} = \mathbb{E}_{X_0^\mu} \int_{\mathcal{X}} \|\nabla \tilde{f}_k^*(x) - \nabla f_k^*(x)\|_2^2 d\tilde{\mu}_k(x) \quad (20)$$

as the (expected) gradient error. In general, it is hard to derive upper bounds of gradients for general functions. Nevertheless, we upper bound function gradients by function values for a specific function class,  $\tilde{\mathcal{F}}$  defined in Section 3.1. Below we provide a generic assumption on  $\tilde{\mathcal{F}}$  to derive the desired upper bounds of gradients.

**Assumption 7** *The set  $\nabla \tilde{\mathcal{F}} = \{\nabla f : f \in \tilde{\mathcal{F}}\}$  is closed, bounded in  $(\mathcal{C}(\mathcal{X}), \ell_\infty)$ . For each  $\nabla f \in \nabla \tilde{\mathcal{F}}$ ,  $\nabla f$  is  $h$ -Lipschitz for some  $h > 0$ .*

Such an assumption can be achieved by function classes with uniformly bounded and Lipschitz continuous gradients, which includes the class of neural networks defined in (15) with bounded parameters. See §D.1 for more details. In what follows, we use Lemma 21 in the appendix to obtain gradient error bounds.

**Generalization error of VFM.** By setting  $p = 2$  and  $f(x) = \tilde{f}_k^*(x) - f_k^*(x)$  in Lemma 21, we are able to upper bound the gradient errors by the generalization errors of NNs, which is bounded in §D.3 with the orders of

$$\bar{\varepsilon}_\mu = \mathcal{O}(N_\mu^{-1/2}), \quad \bar{\varepsilon}_\nu = \mathcal{O}(N_\nu^{-1/2}) \quad (21)$$

by wide enough NNs for player  $\mu$  and player  $\nu$ , respectively. Here  $N_\mu$  and  $N_\nu$  are the numbers of particles for player  $\mu$  and player  $\nu$ , respectively. Such results are standard for the stochastic gradient descent (SGD) over neural networks, since the number of iterations  $t$  in Algorithm 3 is also the sample size  $N_\mu$  (resp.  $N_\nu$ ) in our algorithm.

## 4.2. Convergence to the IFNE for PL Games

Recall that  $G(\mu) = \max_{\nu \in \mathcal{M}} F(\mu, \nu)$ . We define  $L_G = \max_{\mu \in \mathcal{M}} \|\text{grad } G(\mu)\|_{\mu}$ , which is upper bounded since  $G$  is Lipschitz (Lemma E.2) on a compact domain  $\mathcal{M}$  (Proposition 30). We assume that there exist constants  $M_H > 0$  and  $M_G > 0$  such that  $M_H = \max_{\mu, \nu_0 \in \mathcal{M}} [G(\mu) - F(\mu, \nu_0)]$  and  $M_G = \max_{\mu_0 \in \mathcal{M}} G(\mu_0) - G(\mu^*)$ , where  $\mu^* \in \text{argmin}_{\mu \in \mathcal{M}} G(\mu)$ . Under Assumption 3 for  $\xi$ -PL games, we characterize the following sublinear rate to find an IFNE defined in Definition 4 by VTIG with sample sizes  $N_{\mu}, N_{\nu}$  and numbers of iterations  $K_{\mu}, K_{\nu}$ . Recall that  $L_0, L_1$ , and  $L_2$  are Lipschitz constants defined in Assumption 2. The constant  $\xi$  for the PL condition is defined in Assumption 3. Also,  $\sigma = 1 - \xi\eta^{\nu}/2 \in (0, 1)$  is a contraction factor from Lemma 34.

**Theorem 8 (Convergence of Infinite-Dimensional PL Games)** *Suppose that the objective  $F$  admits a variational form under Assumption 2 and 3. Also, the function class  $\tilde{\mathcal{F}}$  satisfies Assumption 7. We set the stepsizes to be  $\eta^{\mu} \in [0, \min\{1/h, 2/\tilde{L}\})$  and  $\eta^{\nu} \in (0, \min\{1/(4L_{\nu}), 1/h\})$ , where  $\tilde{L} = L_1 + L_0^2/\xi$ . Then, for any  $\theta > 0$ , if*

$$K_{\nu} \geq K_{\nu}(\theta) = \mathcal{O}\left(\log\left(\frac{(1-\sigma)\widehat{M}_H - \eta^{\nu}\bar{\varepsilon}_{\nu}}{\theta}\right) / \log\frac{1}{\sigma}\right), \quad \text{where } \widehat{M}_H = \max\left\{M_H, \frac{\eta^{\nu}\bar{\varepsilon}_{\nu} + 1}{1-\sigma}\right\}, \quad (22)$$

there exists an iteration  $k \in [K_{\mu}]$  such that

$$\mathbb{E}_{X_0}[\mathcal{J}_{\mu}^2(\tilde{\mu}_k, \tilde{\nu}_{k+1})] = \mathcal{O}\left((\Delta + \sqrt{\bar{\varepsilon}_{\mu}})^2 \cdot \left((\Delta + \sqrt{\bar{\varepsilon}_{\mu}}) + \frac{M_G}{K_{\mu}}\right)\right), \quad \mathbb{E}_{X_0}[\mathcal{J}_{\nu}(\tilde{\mu}_k, \tilde{\nu}_{k+1})] = \mathcal{O}\left(\frac{L_2\Delta}{L_0}\right). \quad (23)$$

Here  $\Delta = L_0\sqrt{\frac{\eta_{\nu}\bar{\varepsilon}_{\nu} + \theta}{2\xi(1-\sigma)}}$ , and the gradient error terms  $\bar{\varepsilon}_{\mu}$  and  $\bar{\varepsilon}_{\nu}$  are characterized in (21).

**Proof** See §E.4 for a detailed proof and more dependencies on other constants. ■

The proof of Theorem 8 is based on Lemma 34 and a Danskin-type lemma in §E.2 which ensures an appropriate estimate of  $\text{grad } G$  provided by inner loops and the smoothness of the objective defined in Assumption 2. Such properties imply that VTIG behaves as the gradient descent over the inner maximization value functional  $G$ , which concludes the proof. The bounds for the first-order errors  $\mathcal{J}_{\mu}$  and  $\mathcal{J}_{\nu}$  are composed of the optimization error  $\theta$  of player  $\nu$ , the optimization error  $\mathcal{O}(K_{\mu}^{-1})$  of player  $\mu$ , and the gradient errors  $\bar{\varepsilon}_{\mu}$  and  $\bar{\varepsilon}_{\nu}$  characterized in (21) due to finite samples. Specifically, the term  $\Delta$  encapsulates both the statistical error and the optimization error of player  $\nu$ , which are added to  $\sqrt{\bar{\varepsilon}_{\mu}}$  and  $\mathcal{O}(K_{\mu}^{-1})$  in the error bound for player  $\mu$ . Considering  $N_{\mu}, N_{\nu}, K_{\mu}$ , and  $K_{\nu}$  as dominating terms in the bounds, if we set  $N_{\mu} = N_{\nu} = \mathcal{O}(\theta^{-4})$ ,  $K_{\mu} \geq K_{\mu}(\theta) = \mathcal{O}(\theta^{-2})$ , and  $K_{\nu} \geq K_{\nu}(\theta) = \mathcal{O}(\log(\theta^{-1}))$ , by Definition 4 we achieve a  $\theta$ -IFNE. In this sense, VTIG converges at a sublinear rate to the IFNE defined in (10) under the PL game condition.

## 4.3. Convergence to the Minimax Value under the Two-Sided PL Condition

In this section, we aim to achieve a stronger convergence result leading to the minimax value of the game by a stronger assumption. We give the definition of two-sided Riemannian PL games below.

**Assumption 9 (Two-Sided Riemannian PL Game)** *We define functionals  $H_{\mu}(\nu) = -F(\mu, \nu)$  and  $F_{\nu}(\mu) = F(\mu, \nu)$  for fixed  $\mu$  and  $\nu$ , respectively. We assume (3) to be a two-sided Riemannian PL game, or simply a two-sided PL game, in the sense that  $F_{\nu}(\mu)$  is  $\xi_1$ -PL and  $H_{\mu}(\nu)$  is  $\xi_2$ -PL for some  $\xi_1, \xi_2 > 0$ .*

Note that the definition of two-sided PL games relaxes that of the convex-concave games even in infinite-dimensional spaces. In fact, Example 1 provides a two-sided PL game by KL regularization for both players, which is ubiquitous in training GANs. Assumption 9 also guarantees a PL condition on  $G(\mu) = \max_{\nu \in \mathcal{M}} F(\mu, \nu)$  according to Lemma 36. By using such a landscape, we establish the linear convergence rate of finding the minimax value of the game as below.

**Theorem 10 (Convergence to the Minimax Value of Two-Sided PL Games)** *Let the objective  $F$  satisfy (4), Assumption 2 and 9. Suppose that  $\tilde{F}$  satisfies Assumption 7. With the outer-loop stepsize  $\eta^\mu \in [0, \min\{1/h, 1/(4\tilde{L})\}]$  and inner-loop stepsize  $\eta^\nu \in (0, \min\{1/(4L_\nu), 1/h\})$ , for  $k \geq 1$  it holds that*

$$\mathbb{E}_{X_0} \left[ F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)) \right] - F(\mu^*, \nu^*) \leq \underbrace{\tilde{\sigma}^k \cdot \left( \mathbb{E}_{X_0} [F(\tilde{\mu}_0, \tilde{\nu}_1)] - F(\mu^*, \nu^*) \right)}_{(i)} + \underbrace{\frac{1 - \tilde{\sigma}^k}{1 - \tilde{\sigma}} \cdot \eta^\mu (\bar{\varepsilon}_\mu + \tilde{\Delta}^2)}_{(ii)}, \quad (24)$$

where  $\tilde{\mu}_k$  and  $\tilde{\nu}_{k+1}$  are probability measure iterates defined in Algorithm 2, gradient error terms  $\bar{\varepsilon}_\mu$  and  $\bar{\varepsilon}_\nu$  are given in (21). The expectation is taken w.r.t. the initial sample  $X_0$ . The contraction factor is  $\tilde{\sigma} = 1 - \xi_1 \eta^\mu / 2$ , and we define the total error term for player  $\nu$  as  $\tilde{\Delta}^2 = L_0^2 / 2\xi_2 \cdot \left( \sigma^{K_\nu} \cdot M_H + \eta^\nu \bar{\varepsilon}_\nu \cdot \frac{1 - \sigma^{K_\nu}}{1 - \sigma} \right)$ , where  $M_H$  is the upper bound of  $F(\mu, \nu^*(\mu)) - F(\mu, \nu_0(\mu))$  defined in §4.2, and  $K_\nu$  denotes the number of timesteps for player  $\nu$  in Algorithm 2.

**Proof** See §F.2 for a detailed proof. ■

The proof of Theorem 10 differs from that of Theorem 8 mainly by the lower bounds of gradient norms provided by  $\xi_1$ -PL condition on functional  $G$ . Under the two-sided PL condition in Assumption 9, Theorem 10 characterizes a linear convergence rate for VTIG of the objective functional value to the minimax value  $F(\mu^*, \nu^*)$  of the game, with an accumulated statistical error term (ii). In detail, the optimization error (i) decays by a factor of  $\tilde{\sigma}$  linearly. Additionally, our statistical error is composed of the gradient error  $\bar{\varepsilon}_\mu$  of player  $\mu$  and the error term  $\tilde{\Delta}^2$ , which is further decomposed into the linearly decaying optimization error  $\sigma^{K_\nu} M_H$  and the gradient error  $\bar{\varepsilon}_\nu$  of player  $\nu$  scaled by  $(1 - \sigma^{K_\nu}) / (1 - \sigma)$ . Specifically, in the total bound (24)  $\bar{\varepsilon}_\mu$  scales at a rate of  $(1 - \tilde{\sigma}^k) / (1 - \tilde{\sigma})$  with the iteration  $k$  and  $\bar{\varepsilon}_\nu$  scales at the rate of  $(1 - \tilde{\sigma}^k) / (1 - \tilde{\sigma}) \cdot (1 - \sigma^{K_\nu}) / (1 - \sigma)$ , which implies the error accumulation from the inner loop of Algorithm 2. Also, we adopt the objective value instead of IFNE in Theorem 8 to measure the error of convergence to the minimax value. Although we suffer from the finite-sample error to approximate probability measures, it is flexible to tune parameters  $N_\mu, N_\nu, K_\mu$ , and  $K_\nu$  according to their corresponding error terms in the bound to optimize the algorithm in practice, especially when some parameters are restricted.

## Acknowledgement

We thank Baptiste Goujaud and Damien Scieur for enlightening discussion.

## References

- [1] Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein gan. *arXiv preprint arXiv:1701.07875*, 2017.
- [2] Sanjeev Arora, Rong Ge, Yingyu Liang, Tengyu Ma, and Yi Zhang. Generalization and equilibrium in generative adversarial nets (gans). In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 224–232. JMLR. org, 2017.
- [3] Maximilian Balandat, Walid Krichene, Claire J. Tomlin, and Alexandre M. Bayen. Minimizing regret on reflexive banach spaces and learning nash equilibria in continuous zero-sum games. *CoRR*, abs/1606.01261, 2016.
- [4] Fabrice Baudoin and Michel Bonnefont. Reverse poincaré inequalities, isoperimetry, and riesz transforms in carnot groups. *Nonlinear Analysis*, 131:48–59, 2016.
- [5] Pierre Bernhard and Alain Rapaport. On a theorem of danskin with an application to a theorem of von neumann-sion. *Nonlinear Analysis: Theory, Methods & Applications*, 24(8):1163–1181, 1995.
- [6] Dmitri Burago, Sergei Ivanov, and Yuri Burago. Course in metric geometry. 2001.
- [7] Eric A Carlen and Wilfrid Gangbo. Constrained steepest descent in the 2-Wasserstein metric. *Annals of mathematics*, pages 807–846, 2003.
- [8] Antonin Chambolle and Thomas Pock. On the ergodic convergence rates of a first-order primal–dual algorithm. *Mathematical Programming*, 159(1-2):253–287, 2016.
- [9] Changyou Chen, Ruiyi Zhang, Wenlin Wang, Bai Li, and Liqun Chen. A unified particle-optimization framework for scalable bayesian sampling. *arXiv preprint arXiv:1805.11659*, 2018.
- [10] Shiing-Shen Chern, Wei-huan Chen, and Kai Shue Lam. *Lectures on differential geometry*, volume 1. World Scientific Publishing Company, 1999.
- [11] Neil E Cotter. The stone-weierstrass theorem and its application to neural networks. *IEEE Transactions on Neural Networks*, 1(4):290–295, 1990.
- [12] Balázs Csanád Csáji et al. Approximation with artificial neural networks. *Faculty of Sciences, Etvos Lornd University, Hungary*, 24(48):7, 2001.
- [13] Bo Dai, Albert Shaw, Lihong Li, Lin Xiao, Niao He, Zhen Liu, Jianshu Chen, and Le Song. Sbeed: Convergent reinforcement learning with nonlinear function approximation. *arXiv preprint arXiv:1712.10285*, 2017.
- [14] Bo Dai, Hanjun Dai, Arthur Gretton, Le Song, Dale Schuurmans, and Niao He. Kernel exponential family estimation via doubly dual embedding. *arXiv preprint arXiv:1811.02228*, 2018.
- [15] Cong D Dang and Guanghui Lan. On the convergence properties of non-euclidean extragradient methods for variational inequalities with generalized monotone operators. *Computational Optimization and applications*, 60(2):277–310, 2015.
- [16] Constantinos Daskalakis and Ioannis Panageas. Last-iterate convergence: Zero-sum games and constrained min-max optimization. *arXiv preprint arXiv:1807.04252*, 2018.

- [17] Constantinos Daskalakis and Ioannis Panageas. The limit points of (optimistic) gradient descent in min-max optimization. In *Advances in Neural Information Processing Systems*, pages 9236–9246, 2018.
- [18] L.C. Evans. *Partial Differential Equations*. American Mathematical Society, 2010.
- [19] Francisco Facchinei and Jong-Shi Pang. *Finite-dimensional variational inequalities and complementarity problems*. Springer Science & Business Media, 2007.
- [20] Charlie Frogner and Tomaso Poggio. Approximate inference with wasserstein gradient flows. *arXiv preprint arXiv:1806.04542*, 2018.
- [21] A Gaivoronski. Linearization methods for optimization of functionals which depend on probability measures. In *Stochastic Programming 84 Part II*, pages 157–181. Springer, 1986.
- [22] Soumyadip Ghosh, Mark Squillante, and Ebisa Wollega. Efficient stochastic gradient descent for distributionally robust learning. *arXiv preprint arXiv:1805.08728*, 2018.
- [23] Gauthier Gidel, Hugo Berard, Gaëtan Vignoud, Pascal Vincent, and Simon Lacoste-Julien. A variational inequality perspective on generative adversarial networks. *arXiv preprint arXiv:1802.10551*, 2018.
- [24] Gauthier Gidel, Reyhane Askari Hemmat, Mohammad Pezeshki, Remi Lepriol, Gabriel Huang, Simon Lacoste-Julien, and Ioannis Mitliagkas. Negative momentum for improved game dynamics. *arXiv preprint arXiv:1807.04740*, 2018.
- [25] Nicola Gigli. On the inverse implication of brenier-mccann theorems and the structure of  $(p_2(\mathcal{M}), w_2)$ . *Methods and Applications of Analysis*, 18(2):127–158, 2011.
- [26] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In *Advances in neural information processing systems*, pages 2672–2680, 2014.
- [27] Paulina Grnarova, Kfir Y Levy, Aurelien Lucchi, Thomas Hofmann, and Andreas Krause. An online learning approach to generative adversarial networks. *arXiv preprint arXiv:1706.03269*, 2017.
- [28] E Yazdandoost Hamedani, A Jalilzadeh, NS Aybat, and UV Shanbhag. Iteration complexity of randomized primal-dual methods for convex-concave saddle point problems. *arXiv preprint arXiv:1806.04118*, 2018.
- [29] Jonathan Ho and Stefano Ermon. Generative adversarial imitation learning. In *Advances in neural information processing systems*, pages 4565–4573, 2016.
- [30] Thomas Hofmann, Bernhard Schölkopf, and Alexander J Smola. Kernel methods in machine learning. *Annals of Statistics*, pages 1171–1220, 2008.
- [31] Ya-Ping Hsieh, Chen Liu, and Volkan Cevher. Finding mixed nash equilibria of generative adversarial networks. *arXiv preprint arXiv:1811.02002*, 2018.
- [32] Chi Jin, Praneeth Netrapalli, and Michael I Jordan. Minmax optimization: Stable limit points of gradient descent ascent are locally optimal. *arXiv preprint arXiv:1902.00618*, 2019.

- [33] Anatoli Juditsky and Arkadi Nemirovski. Solving variational inequalities with monotone operators on domains given by linear minimization oracles. *Mathematical Programming*, 156(1-2):221–256, 2016.
- [34] Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- [35] John M Lee. *Introduction to smooth manifolds*. Springer, 2001.
- [36] Chang Liu, Jingwei Zhuo, Pengyu Cheng, Ruiyi Zhang, Jun Zhu, and Lawrence Carin. Understanding and accelerating particle-based variational inference. *arXiv preprint arXiv:1807.01750*, 2018.
- [37] Yuanyuan Liu, Fanhua Shang, James Cheng, Hong Cheng, and Licheng Jiao. Accelerated first-order methods for geodesically convex optimization on riemannian manifolds. In *Advances in Neural Information Processing Systems*, pages 4868–4877, 2017.
- [38] Songtao Lu, Ioannis Tsaknakis, and Mingyi Hong. Block alternating optimization for non-convex min-max problems: algorithms and applications in signal processing and communications. In *ICASSP 2019-2019 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 4754–4758. IEEE, 2019.
- [39] Songtao Lu, Ioannis Tsaknakis, Mingyi Hong, and Yongxin Chen. Hybrid block successive approximation for one-sided non-convex min-max problems: algorithms and applications. *arXiv preprint arXiv:1902.08294*, 2019.
- [40] Tung Mai, Milena Mihail, Ioannis Panageas, Will Ratcliff, Vijay Vazirani, and Peter Yunker. Cycles in zero-sum differential games and biological diversity. In *Proceedings of the 2018 ACM Conference on Economics and Computation*, pages 339–350, 2018.
- [41] Lars Mescheder, Andreas Geiger, and Sebastian Nowozin. Which training methods for gans do actually converge? *arXiv preprint arXiv:1801.04406*, 2018.
- [42] Ilya Molchanov and Sergei Zuyev. Variational calculus in the space of measures and optimal design. In *Optimum design 2000*, pages 79–90. Springer, 2001.
- [43] Renato DC Monteiro and Benar Fux Svaiter. On the complexity of the hybrid proximal extragradient method for the iterates and the ergodic mean. *SIAM Journal on Optimization*, 20(6):2755–2787, 2010.
- [44] Arkadi Nemirovski. Prox-method with rate of convergence  $o(1/t)$  for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15(1):229–251, 2004.
- [45] Yurii Nesterov and Boris T Polyak. Cubic regularization of newton method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.
- [46] XuanLong Nguyen, Martin J Wainwright, and Michael I Jordan. Estimating divergence functionals and the likelihood ratio by convex risk minimization. *IEEE Transactions on Information Theory*, 56(11): 5847–5861, 2010.

- [47] Sebastian Nowozin, Botond Cseke, and Ryota Tomioka. f-gan: Training generative neural samplers using variational divergence minimization. In *Advances in neural information processing systems*, pages 271–279, 2016.
- [48] Frans A Oliehoek, Rahul Savani, Jose Gallego, Elise van der Pol, and Roderich Groß. Beyond local nash equilibria for adversarial networks. In *Benelux Conference on Artificial Intelligence*, pages 73–89. Springer, 2018.
- [49] Felix Otto and Cédric Villani. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000.
- [50] Jong-Shi Pang and Meisam Razaviyayn. A unified distributed algorithm for non-cooperative games., 2016.
- [51] Peter Petersen, S Axler, and KA Ribet. *Riemannian geometry*, volume 171. Springer, 2006.
- [52] Boris Teodorovich Polyak. Gradient methods for minimizing functionals. *Zhurnal Vychislitel’noi Matematiki i Matematicheskoi Fiziki*, 3(4):643–653, 1963.
- [53] Maxim Raginsky, Alexander Rakhlin, and Matus Telgarsky. Non-convex learning via stochastic gradient langevin dynamics: a nonasymptotic analysis. *arXiv preprint arXiv:1702.03849*, 2017.
- [54] Pierre H Richemond and Brendan Maginnis. On wasserstein reinforcement learning and the fokker-planck equation. *arXiv preprint arXiv:1712.07185*, 2017.
- [55] R Tyrrell Rockafellar. *Convex analysis*. Number 28. Princeton university press, 1970.
- [56] Walter Rudin. Principles of mathematical analysis. 1976.
- [57] Maziar Sanjabi, Jimmy Ba, Meisam Razaviyayn, and Jason D Lee. On the convergence and robustness of training gans with regularized optimal transport. In *Advances in Neural Information Processing Systems*, pages 7091–7101, 2018.
- [58] Maziar Sanjabi, Meisam Razaviyayn, and Jason D Lee. Solving non-convex non-concave min-max games under polyak-Łojasiewicz condition. *arXiv preprint arXiv:1812.02878*, 2018.
- [59] Aman Sinha, Hongseok Namkoong, and John Duchi. Certifying some distributional robustness with principled adversarial training. *arXiv preprint arXiv:1710.10571*, 2017.
- [60] J Stewart. Chapter 15.2 limits and continuity, multivariable calculus, 2008.
- [61] Kiran K Thekumparampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Efficient algorithms for smooth minimax optimization. In *Advances in Neural Information Processing Systems*, pages 12659–12670, 2019.
- [62] Bart PG Van Parys, Peyman Mohajerin Esfahani, and Daniel Kuhn. From data to decisions: Distributionally robust optimization is optimal. *arXiv preprint arXiv:1704.04118*, 2017.
- [63] Cédric Villani. *Topics in optimal transportation*. Number 58. American Mathematical Society, 2003.
- [64] Cédric Villani. *Optimal transport: old and new*, volume 338. Springer, 2008.

- [65] Andre Wibisono. Sampling as optimization in the space of measures: The langevin dynamics as a composite optimization problem. *arXiv preprint arXiv:1802.08089*, 2018.
- [66] Hongyi Zhang and Suvrit Sra. An estimate sequence for geodesically convex optimization. In *Conference On Learning Theory*, pages 1703–1723, 2018.
- [67] Hongyi Zhang, Sashank J Reddi, and Suvrit Sra. Riemannian svrg: Fast stochastic optimization on riemannian manifolds. In *Advances in Neural Information Processing Systems*, pages 4592–4600, 2016.
- [68] Difan Zou, Pan Xu, and Quanquan Gu. Subsampled stochastic variance-reduced gradient langevin dynamics. In *International Conference on Uncertainty in Artificial Intelligence*, 2018.



## Appendix A. Additional Algorithms

---

**Algorithm 2** Theoretical Version of Multi-Step Variational Transport Algorithm for Infinite-Dimensional Games (VTIG) with Resampling

---

- 1: **Input:** Functional  $F: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$ ; initial measures  $\tilde{\mu}_0 \in \mathcal{M}, \tilde{\nu}_0 \in \mathcal{M}$ ; numbers of particles  $N_\mu, N_\nu$ ; numbers of iterations  $K_\mu, K_\nu$ ; and stepsizes  $\eta_k^\mu \in (0, \min\{1/h, 2/\tilde{L}\})$ ,  $\eta^\nu \in (0, \min\{1/(4L_2), 1/h\})$ .
  - 2: Initialize the transportation maps  $T_0^\mu \leftarrow \text{id}, T_0^\nu \leftarrow \text{id}$ .
  - 3: Initialize  $N_\mu$  particles  $X_0^\mu = \{x_i^\mu\}_{i \in [N_\mu]}$  by drawing  $N_\mu$  i.i.d. samples from  $\tilde{\mu}_0$ .
  - 4: Initialize  $N_\nu$  particles  $X_0^\nu = \{x_i^\nu\}_{i \in [N_\nu]}$  by drawing  $N_\nu$  i.i.d. samples from  $\tilde{\nu}_0$ .
  - 5: **for**  $k = 0, 1, 2, \dots, K_\mu - 1$  **do**
  - 6:   Generate  $N_\mu$  particles  $X_0^\mu = \{x_i^\mu\}_{i \in [N_\mu]}$  by drawing  $N_\mu$  i.i.d. samples from  $\tilde{\mu}_0$ .
  - 7:   Push  $\mu$ -particles by letting  $X_k^\mu \leftarrow T_k^\mu(X_0^\mu)$ .
  - 8:   **for**  $l = 0, 1, 2, \dots, K_\nu - 1$  **do**
  - 9:     Generate  $N_\nu$  particles  $X_0^\nu(\tilde{\mu}_k) = \{x_i^\nu\}_{i \in [N_\nu]}$  by drawing  $N_\nu$  i.i.d. samples from  $\tilde{\nu}_0$ .
  - 10:     Push  $\nu$ -particles by letting  $X_l^\nu(\tilde{\mu}_k) \leftarrow T_{kK_\nu+l}^\nu(X_0^\nu(\tilde{\mu}_k))$ .
  - 11:      $\tilde{f}_{l,k}^* \leftarrow \mathbf{VFM}(X_l^\nu(\tilde{\mu}_k), F_{\tilde{\mu}_k}^*, N_\nu)$ .
  - 12:     Update the transportation map by letting  $T_{kK_\nu+l+1}^\nu = [\text{Exp}_{\mathcal{X}}(-\eta^\nu \cdot \nabla \tilde{f}_{l,k}^*)] \circ T_{kK_\nu+l}^\nu$ .
  - 13:   **end for**
  - 14:    $\tilde{f}_k^* \leftarrow \mathbf{VFM}(X_k^\mu, F_{\tilde{\nu}_{k+1}}^*, N_\mu)$ .
  - 15:   Update the transportation map by letting  $T_{k+1}^\mu = [\text{Exp}_{\mathcal{X}}(-\eta_k^\mu \cdot \nabla \tilde{f}_k^*)] \circ T_k^\mu$ .
  - 16: **end for**
  - 17: **Output:** The final transportation maps  $T_{K_\mu}^\mu$  and  $T_{K_\mu K_\nu}^\nu$ .
- 

---

**Algorithm 3** Variational Form Maximization via SGD ( $\mathbf{VFM}(\{x_i\}_{i=1}^t, F^*, t)$ )

---

- 1: **Require:** Initial weights  $b_i, [\beta(0)]_i$  ( $i \in [w]$ ), number of iterations  $t$ , sample  $\{x_i\}_{i=1}^t$ , and projection radius  $r_f$ .
  - 2: Set stepsize  $\eta \leftarrow t^{-1/2}$
  - 3: **for**  $s = 0, \dots, t - 1$  **do**
  - 4:    $x \leftarrow x_{s+1}$
  - 5:    $\beta(s + 1/2) \leftarrow \beta(s) - \eta (\nabla_\beta F^*(f_{\beta(s)}(x)) - \nabla_\beta f_{\beta(s)}(x))$
  - 6:    $\beta(s + 1) \leftarrow \text{argmin}_{\beta \in \mathcal{B}^0(r_f)} \{\|\beta - \beta(s + 1/2)\|_2\}$
  - 7: **end for**
  - 8: Average over path  $\hat{\beta} \leftarrow 1/t \cdot \sum_{s=0}^{t-1} \beta(s)$
  - 9: **Output:**  $f_{\hat{\beta}}$
- 

Algorithm 2 provide the resampling version of VTIG (Algorithm 1) for theoretical analysis in §4. As mentioned above, here we explicitly use two sequences of transportation maps  $\{T_k^\mu\}_{k=0}^{K_\mu}$  and  $\{T_k^\nu\}_{k=0}^{K_\mu K_\nu}$  to record the transportation plans of mapping the initial probability measures to the current iterates of Player  $\mu$  and player  $\nu$ , respectively. For simplicity, we put the particle pushing step before the VFM step, while the essential difference between Algorithm 2 and Algorithm 1 lies only in the resampling procedures at the beginning of each player's loop, *i.e.*, Line 5 and Line 8 in Algorithm 2, after which we adopt the recorded transportation plan from time-step 0 to the current timestep  $k$ ,  $T_k^\mu$  or  $T_k^\nu$ , to push the resampled initial particles into their states at the current timestep through  $k$  steps of transportation maps. In contrast, in Algorithm 1 we push the two sets of particles for one step at each timestep based on their current states, with

only one sampling procedure before the beginning of the two loops. Hence, We remark that Algorithm 1 is deterministic in essence; the only randomness comes from the initialization of the particles. In addition, note that even in resampling cases, our algorithm only requires to call an oracle that is able to sample from a fixed distribution, which is more efficient than SGLD requiring Markov-chain type sampling for different probability measure iterates. We remark that Algorithm 2 is considered only for the sake of theoretical analysis; random sampling is unnecessary in practical implementation.

We also remark that the exponential maps  $\text{Exp}_{\mathcal{X}}(-\eta^\mu \cdot \nabla \tilde{f}_k^*)$  in Algorithm 1 and 2 are essentially translations of particles in the direction of  $-\nabla \tilde{f}_k^*$  with a stepsize  $\eta^\nu$  in  $\mathcal{X}$ , which implies that they are computationally efficient for infinite-dimensional problems.

## Appendix B. Properties of Optimization over Riemannian Manifolds

### B.1. Background Details

In this section, we present basic properties of functions and gradient descent on a Riemannian manifold  $(\mathcal{M}, g)$  with a Riemannian metric  $g$ , supposing that any two points on  $\mathcal{M}$  uniquely determines a geodesic. By the nature of Wasserstein spaces, these propositions also apply to our optimization process over the family of distributions. With such background, we are better prepared to demonstrate the convergence results by extending the classical optimization arguments to Riemannian manifolds. For further extensions of definitions and propositions within this section, we refer to other literatures on geodesically convex optimization. See, e.g., [37, 66, 67] and the references therein.

### B.2. Riemannian Manifold

Let  $\mathcal{M}$  be a  $d$ -dimensional differential manifold, *i.e.*, a topological space that is locally homeomorphic to the Euclidean space  $\mathbb{R}^d$  and has a globally defined differential structure [10].

**Definition 11 (Tangent vector)** *A tangent vector at  $x \in \mathcal{M}$  is an equivalence class of differentiable curves through  $x$  with a prescribed velocity vector at  $x$ . The tangent space at  $x$ , denoted by  $T_x\mathcal{M}$ , consists of all tangent vectors at  $x$ .*

In what follows, we denote by  $f : \mathcal{M} \rightarrow \mathbb{R}$  a differentiable function over  $\mathcal{M}$ , and define its directional derivative as  $\nabla_v f(x) = \frac{d}{dt} f[\gamma(t)]|_{t=0}$ , where  $\gamma$  is any smooth curve such that  $\gamma(0) = x$  and  $\gamma'(0) = v$ .

To compare two tangent vectors in a meaningful way, we consider a Riemannian manifold  $(\mathcal{M}, g)$ , which is a real smooth manifold equipped with an Riemannian metric  $g_x$  on the tangent space  $T_x\mathcal{M}$  for any  $x \in \mathcal{M}$  [51]. The inner product of any two tangent vectors  $u_1, u_2 \in T_x\mathcal{M}$  is defined as  $\langle u_1, u_2 \rangle_x = g_x(u_1, u_2)$ , with an induced norm  $\|u_1\|_x = \sqrt{\langle u_1, u_1 \rangle_x}$ . We specify how to move a point along a direction in the following definition.

**Definition 12 (Exponential map)** *The exponential map at  $x$ , denoted by  $\text{Exp}_x$ , sends any tangent vector  $u \in T_x\mathcal{M}$  to  $y = \gamma_u(1) \in \mathcal{M}$ , where  $\gamma_u : [0, 1] \rightarrow \mathcal{M}$  is the unique geodesic determined by  $\gamma_u(0) = x$  and  $\gamma_u'(0) = u$ .*

Moreover, since  $\gamma_u$  is the unique geodesic connecting  $x$  and  $y$ , the exponential map is invertible and we have  $u = \text{Exp}_x^{-1}(y)$ . The distance between  $x$  and  $y$  satisfies  $d(x, y) = [\langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(y) \rangle_x]^{1/2}$ , which is also called the geodesic distance. For any two points  $x, y \in \mathcal{M}$ , the parallel transport  $\Gamma_x^y : T_x\mathcal{M} \rightarrow T_y\mathcal{M}$  specifies that how a tangent vector of  $x$  is identified with an element in  $T_y\mathcal{M}$ . Moreover, we have  $\langle u, v \rangle_x = \langle \Gamma_x^y u, \Gamma(\gamma)_{xv}^y \rangle_y$  for any  $u, v \in T_x\mathcal{M}$ . For simplicity, we denote  $d^2(u, v) = \langle u - \Gamma_y^x v, u - \Gamma_y^x v \rangle_x$  for any

$x, y \in \mathcal{M}$  and any  $u \in T_x\mathcal{M}, v \in T_y\mathcal{M}$ . Below we define the gradient *w.r.t.* the Riemannian metric, in contrast with the Euclidean space.

**Definition 13 (Gradient)** Suppose there exists  $u \in T_x\mathcal{M}$  such that  $\nabla_v f(x) = \langle u, v \rangle_x$  for any  $v \in T_x\mathcal{M}$ , then  $f$  is differentiable at  $x$ , and  $u$  is called the gradient of  $f$  at  $x$ , denoted by  $\text{grad } f(x)$ . We also define partial gradient of  $f$  at  $(x, y)$  w.r.t.  $x$  by  $\text{grad}_x f(x, y) = \text{grad } f_y(x)$  as a natural extension, where  $f_y(x) = f(x, y)$  for any fixed  $y \in \mathcal{M}$ . We call function  $f$   $W_2$ -differentiable if  $\text{grad } f$  exists over its domain.

**Definition 14 (Geodesic space)** A metric space  $(\mathcal{X}, d)$  consists of a set  $\mathcal{X}$  and a distance function  $d: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  satisfying a few simple properties [6]. A curve  $\gamma$  on  $\mathcal{X}$  is a continuous function from  $[0, 1]$  to  $\mathcal{X}$ , whose length is defined as  $L(\gamma) = \sup \sum_{i=1}^n d[\gamma(t_{i-1}), \gamma(t_i)]$ , where the supremum is taken over  $n \geq 1$  and all partitions  $0 = t_0 < t_1 < \dots < t_n = 1$  of  $[0, 1]$ . By this definition, for any curve  $\gamma$ , we have  $L(\gamma) \geq d[\gamma(0), \gamma(1)]$ . If there exists a constant  $v \geq 0$  such that,  $d[\gamma(t_1), \gamma(t_2)] = v \cdot |t_1 - t_2|$  for any  $t_1, t_2 \in [0, 1]$ , then curve  $\gamma$  is called a geodesic. In this case, for any  $0 \leq t_1 < t_2 \leq 1$ , the length of  $\gamma$  restricted to  $[t_1, t_2]$  is equal to  $d[\gamma(t_1), \gamma(t_2)]$ . Thus, a geodesic is everywhere locally a distance minimizer. Moreover,  $(\mathcal{X}, d)$  is called a geodesic space if any two points  $x, y \in \mathcal{X}$  are connected by a geodesic  $\gamma$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .

**Definition 15 (Geodesic convexity)** A function  $f: \mathcal{M} \rightarrow \mathbb{R}$  is called geodesically convex if for any  $x, y \in \mathcal{M}$  and a geodesic  $\gamma: [0, 1] \rightarrow \mathcal{M}$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ , we have

$$f[\gamma(t)] \leq t \cdot f[\gamma(0)] + (1 - t) \cdot f[\gamma(1)], \quad \forall t \in [0, 1]. \quad (25)$$

The following lemma characterizes the geodesic convexity using the gradient of  $f$ .

**Lemma 16** If  $f: \mathcal{M} \rightarrow \mathbb{R}$  is differentiable, then it is geodesically convex if and only if

$$f(y) \geq f(x) + \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x, \quad \forall x, y \in \mathcal{M}. \quad (26)$$

**Proof** For any  $x, y \in \mathcal{M}$ , let  $\gamma: [0, 1] \rightarrow \mathcal{M}$  be the unique geodesic satisfying  $\gamma(0) = x$  and  $\gamma(1) = y$ . By the definition of exponential map, we have  $\text{Exp}_x[\gamma'(0)] = y$ , i.e.,  $\gamma'(0) = \text{Exp}_x^{-1}(y)$ . In addition, (25) shows that  $f[\gamma(t)]$  is a convex and differentiable function on  $[0, 1]$ , which implies

$$f[\gamma(1)] \geq f[\gamma(0)] + \left. \frac{d}{dt} f[\gamma(t)] \right|_{t=0}. \quad (27)$$

By the definition of directional derivative, we have

$$\left. \frac{d}{dt} f[\gamma(t)] \right|_{t=0} = \nabla_{\gamma'(0)} f(x) = \langle \text{grad } f(x), \gamma'(0) \rangle_x = \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x. \quad (28)$$

Thus, combining (27) and (28) we obtain (26).

It remains to show (25) assuming (26) holds. For any geodesic  $\gamma: [0, 1] \rightarrow \mathcal{M}$ , we will show that  $f[\gamma(t)]$  is convex on  $[0, 1]$ . To see this, for any  $0 \leq t_1 \leq t_2 \leq 1$ , let  $x = \gamma(t_1)$  and  $y = \gamma(t_2)$ . Note that we can reparametrize  $\gamma$  to obtain a new geodesic  $\hat{\gamma}$  with  $\hat{\gamma}(0) = x$  and  $\hat{\gamma}(1) = y$  by letting

$$\hat{\gamma}(t) = \gamma[t_1 + (t_2 - t_1) \cdot t]$$

for any  $t \in [0, 1]$ . Since  $\hat{\gamma}$  is a geodesic, by the definition of exponential map, we have

$$\text{Exp}_x^{-1}(y) = \hat{\gamma}'(0) = (t_2 - t_1) \cdot \gamma'(t_1).$$

Thus, by (26), we have

$$\begin{aligned} f[\gamma(t_2)] &= f(y) \geq f(x) + \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x \\ &= f[\gamma(t_1)] + \langle \text{grad } f[\gamma(t_1)], (t_2 - t_1) \cdot \gamma'(t_1) \rangle_{\gamma(t_1)} = f[\gamma(t_1)] + (t_2 - t_1) \cdot \left. \frac{d}{dt} f[\gamma(t)] \right|_{t=t_1}, \end{aligned}$$

which implies that  $f[\gamma(t)]$  is a convex function on  $[0, 1]$ . Thus, (25) holds and we conclude the proof of this lemma.  $\blacksquare$

In the following, we extend the concepts of strong convexity and smoothness to manifold optimization.

**Definition 17 (Geodesic strong convexity and smoothness)** For any  $\mu > 0$ , a differentiable function  $f: \mathcal{M} \rightarrow \mathbb{R}$  is called geodesically  $\mu$ -strongly convex if

$$f(y) \geq f(x) + \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x + \mu/2 \cdot d^2(x, y),$$

where  $d$  is the distance function induced by the Riemannian metric. Moreover,  $f$  is said to be geodesically  $L$ -smooth if  $\text{grad } f$  is  $L$ -Lipschitz continuous. That is, for any  $x, y \in \mathcal{M}$ , we have

$$\langle \text{grad } f(x) - \Gamma_y^x[\text{grad } f(y)], \text{grad } f(x) - \Gamma_y^x[\text{grad } f(y)] \rangle_x \leq L^2 \cdot d^2(x, y), \quad (29)$$

where  $\Gamma_y^x: \mathcal{T}_y \mathcal{M} \rightarrow \mathcal{T}_x \mathcal{M}$  is the parallel transport from the tangent space at  $y$  to that at  $x$ .

Note that we apply the parallel transport in (29) to compare  $\text{grad } f(x)$  and  $\text{grad } f(y)$ , which belong to two different tangent spaces. In the following, we introduce the notion of gradient dominated function.

**Definition 18 (Gradient dominance)** Let  $\mu > 0$  and  $f: \mathcal{M} \rightarrow \mathbb{R}$  be a differentiable function with  $f^* = \min_{x \in \mathcal{M}} f(x)$ . Then  $f$  is  $\mu$ -gradient dominated if

$$2\mu \cdot [f(x) - f^*] \leq \langle \text{grad } f(x), \text{grad } f(x) \rangle_x, \quad \forall x \in \mathcal{M}. \quad (30)$$

In the following lemma, we show that, similar to functions in the Euclidean space,  $\text{grad } f$  being Lipschitz smooth implies that  $f$  can be upper bounded by the distance function. More importantly, we show that, gradient dominance is implied by geodesically strong convexity and thus is a weaker condition.

**Lemma 19** If  $f: \mathcal{M} \rightarrow \mathbb{R}$  is geodesically  $\mu$ -strongly convex, then  $f$  is also  $\mu$ -gradient dominated. In addition, if  $f$  has  $L$ -Lipschitz continuous gradient, then, we have

$$f(y) \leq f(x) + \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x + L/2 \cdot d^2(x, y), \quad \forall x, y \in \mathcal{M}. \quad (31)$$

**Proof** For the first part, let  $f$  be a geodesically  $\mu$ -strongly convex function. Since  $(\mathcal{M}, g)$  is a geodesic space, we have  $d^2(x, y) = \langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(y) \rangle_x$ . Thus, by direct computation, we have

$$\begin{aligned} f(y) &\geq f(x) + \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x + \mu/2 \cdot \langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(y) \rangle_x \\ &= f(x) + \mu/2 \cdot \left\langle \text{Exp}_x^{-1}(y) + 1/\mu \cdot \text{grad } f(x), \text{Exp}_x^{-1}(y) + 1/\mu \cdot \text{grad } f(x) \right\rangle_x \\ &\quad - 1/(2\mu) \cdot \langle \text{grad } f(x), \text{grad } f(x) \rangle_x \\ &\geq f(x) - 1/(2\mu) \cdot \langle \text{grad } f(x), \text{grad } f(x) \rangle_x. \end{aligned} \quad (32)$$

Setting  $y = x^*$  such that  $f(x^*) = f^*$  in (32), we establish (30).

For the second part of Lemma 19, for any  $x, y \in \mathcal{M}$ , let  $\gamma$  be the unique geodesic satisfying  $\gamma(0) = x$  and  $\gamma(1) = y$ . Then we have  $\text{Exp}_x^{-1}(y) = \gamma'(0)$ . Moreover, for any  $t \in [0, 1]$ , note that  $\gamma'(t) \in \mathcal{T}_{\gamma(t)}\mathcal{M}$ . By the definition of the parallel transport, we have  $\Gamma_{\gamma(t)}^x \gamma'(t) = \gamma'(0) = \text{Exp}_x^{-1}(y)$ . Thus, by (28) it holds that

$$\begin{aligned} f(y) - f(x) - \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x &= f[\gamma(1)] - f[\gamma(0)] - \left. \frac{d}{dt} f[\gamma(t)] \right|_{t=0} \\ &= \int_0^1 \{ \langle \text{grad } f[\gamma(t)], \gamma'(t) \rangle_{\gamma(t)} - \langle \text{grad } f(x), \gamma'(0) \rangle_x \} dt \\ &= \int_0^1 \left( \langle \Gamma_{\gamma(t)}^x \{ \text{grad } f[\gamma(t)] \} - \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x \right) dt, \end{aligned} \quad (33)$$

where in the last equality we transport the tangent vectors to  $\mathcal{T}_x\mathcal{M}$ . Besides, by (29) and Cauchy-Schwarz inequality, for any  $z \in \mathcal{M}$ , we have

$$\begin{aligned} &| \langle \text{grad } f(x) - \Gamma_z^x [\text{grad } f(z)], \text{Exp}_x^{-1}(y) \rangle_x | \\ &\leq \{ \langle \text{grad } f(x) - \Gamma_z^x [\text{grad } f(z)], \text{grad } f(x) - \Gamma_z^x [\text{grad } f(z)] \rangle_x \}^{1/2} \cdot [ \langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(y) \rangle_x ]^{1/2} \\ &\leq L \cdot d(x, y) \cdot d(z, x). \end{aligned} \quad (34)$$

Finally, combining (33) and (34), we have

$$f(y) - f(x) - \langle \text{grad } f(x), \text{Exp}_x^{-1}(y) \rangle_x \leq L \cdot \int_0^1 d(x, y) \cdot d[\gamma(t), x] dt = L/2 \cdot d^2(x, y),$$

where the last equality follows from the fact that  $d[\gamma(t), x] = d[\gamma(t), \gamma(0)] = t \cdot d(x, y)$ . Therefore, we establish (31) and conclude the proof of Lemma 19.  $\blacksquare$

## Appendix C. Auxiliary Results

In this section, we collect a bunch of supportive proofs and concrete examples to characterize gradient descent over Wasserstein spaces by pushing particles in Euclidean spaces and to illustrate the feasibility of the variational form. First, we show that the Riemannian gradient in distribution spaces can be expressed in the functional gradient in vector variable spaces.

### C.1. Proof of Proposition 5

**Proof** We start from the definition of directional derivative to introduce the link between Riemannian gradients and derivatives *w.r.t.*  $\ell_2$ -norm. For any  $s \in T_\mu\mathcal{M}$  and any  $\mu \in \mathcal{M}$  with corresponding density  $p$ , suppose  $\gamma: [0, 1] \rightarrow \mathcal{M}$  represents a curve satisfying  $\gamma(0) = p$  and  $\gamma'(0) = s$ . Then, the directional derivative of  $F$  gives ( Definition 13 )

$$\left. \frac{d}{dt} F[\gamma(t)] \right|_{t=0} = \langle \text{grad } F(\mu), s \rangle_\mu. \quad (35)$$

Furthermore, by the chain rule of functional gradient of  $F$  with respect to the  $\ell_2$ -Euclidean structure, the directional derivative at  $p$  in the direction of  $s$  can be expressed as

$$\left. \frac{d}{dt} F[\gamma(t)] \right|_{t=0} = \int_{\mathcal{X}} \frac{\delta F}{\delta p}(x) \cdot s(x) dx. \quad (36)$$

On the other hand, let  $u: \mathcal{X} \rightarrow \mathbb{R}$  be the unique solution to elliptic equation

$$-\operatorname{div}[p(x) \cdot (\nabla u)(x)] = s(x), \quad \forall x \in \mathcal{X}, \quad (37)$$

where  $\operatorname{div}$  is the divergence operator on  $\mathcal{X}$ . Thus, plugging (36) into (37) we have

$$\begin{aligned} \frac{d}{dt} F[\gamma(t)] \Big|_{t=0} &= - \int_{\mathcal{X}} \frac{\delta F}{\delta p}(x) \cdot \operatorname{div}[p(x) \cdot (\nabla u)(x)] \, dx \\ &= - \int_{\mathcal{X}} \left\{ \operatorname{div} \left[ p(x) \cdot \frac{\delta F}{\delta p}(x) \cdot (\nabla u)(x) \right] - \left\langle \nabla \left( \frac{\delta F}{\delta p} \right)(x), (\nabla u)(x) \right\rangle \cdot p(x) \right\} \, dx, \end{aligned} \quad (38)$$

where the second equation follows from integration by parts and the property of the divergence operator that

$$\operatorname{div}(f \cdot v) = \langle \nabla f, v \rangle + f \cdot \operatorname{div}(v)$$

holds for any scalar function  $f$  and any vector field  $v$ . It is well known that given some regularity condition, one can show that the first term on the right-hand side of (38) vanishes. For example, when  $\mathcal{X}$  is a convex compact region with periodic boundary condition, this is implied by the divergence theorem [56]. Therefore, again combining (35) and (38) we obtain that

$$\langle \operatorname{grad} F(\mu), s \rangle_{\mu} = \int_{\mathcal{X}} \left\langle \nabla \left( \frac{\delta F}{\delta p} \right)(x), (\nabla u)(x) \right\rangle \cdot p(x) \, dx. \quad (39)$$

Meanwhile we rewrite (37) in  $\operatorname{grad} F(\mu)$ , *i.e.*, another tangent vector at  $\mu$ , with  $v: \mathcal{X} \rightarrow \mathbb{R}$  as the solution to elliptic equation

$$-\operatorname{div}[p(x) \cdot (\nabla v)(x)] = [\operatorname{grad} F(\mu)](x), \quad \forall x \in \mathcal{X}.$$

Since both  $\operatorname{grad} F(\mu)$  and  $s$  lie in the same tangent space  $T_{\mu}\mathcal{M}$ , their inner product is characterized by the Riemannian metric on  $(\mathcal{M}, W_2)$ , that is,

$$\langle \operatorname{grad} F(\mu), s \rangle_{\mu} = \int_{\mathcal{X}} \left\langle (\nabla u)(x), (\nabla v)(x) \right\rangle \cdot p(x) \, dx. \quad (40)$$

Now combining (39) and (40) we have that

$$\int_{\mathcal{X}} \left\langle \nabla \left( \frac{\delta F}{\delta p} \right)(x), (\nabla u)(x) \right\rangle \cdot p(x) \, dx = \langle \operatorname{grad} F(p), s \rangle_p = \int_{\mathcal{X}} \left\langle (\nabla u)(x), (\nabla v)(x) \right\rangle \cdot p(x) \, dx \quad (41)$$

holds for any  $s \in T_{\mu}\mathcal{M}$ . Since  $s$  is arbitrarily picked, (41) indicates that  $\nabla(\delta F/\delta p) = \nabla v$ . To conclude, we have

$$\operatorname{grad} F(\mu) = -\operatorname{div}(p \nabla v) = -\operatorname{div} \left[ p \cdot \nabla \left( \frac{\delta F}{\delta p} \right) \right], \quad (42)$$

which implies the desired result of (12).

In what follows, we turn to obtain the explicit form of  $\delta F/\delta p$  for  $F$  defined in (4), while writing  $F$  as a functional  $F(p)$  in  $p$ . Following the definition of the functional derivative by limits with respect to the  $\ell_2$ -Euclidean structure, for any square-integrable function  $\varphi: \mathcal{X} \rightarrow \mathbb{R}$  in  $\mathcal{F}$ , we can write

$$\int_{\mathcal{X}} \frac{\delta F}{\delta p}(x) \cdot \varphi(x) \, dx = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \cdot [F(p + \epsilon \cdot \varphi) - F(p)]. \quad (43)$$

For simplicity of notations, we denote by  $f_{\epsilon}^*$  the optimal dual solution to the optimization problem

$$\sup_{f \in \mathcal{F}} \left\{ \int_{\mathcal{X}} f(x) \cdot [p(x) + \epsilon \cdot \varphi(x)] \, dx - F^*(f) \right\} \quad (44)$$

for any  $\epsilon$  sufficiently small. Then by setting  $f$  to  $f_p^*$  in the definition of the variational form for  $F(p + \epsilon \cdot \varphi)$ , the difference in (43) satisfies the following lower bound,

$$\begin{aligned} & F(p + \epsilon \cdot \varphi) - F(p) \\ & \geq \left[ \int_{\mathcal{X}} f_p^*(x) \cdot [p(x) + \epsilon \cdot \varphi(x)] \, dx - F^*(f_p^*) \right] - \left[ \int_{\mathcal{X}} f_p^*(x) \cdot p(x) \, dx - F^*(f_p^*) \right] \\ & = \epsilon \cdot \int_{\mathcal{X}} f_p^*(x) \cdot \varphi(x) \, dx. \end{aligned} \quad (45)$$

Meanwhile, we obtain an upper bound of  $F(p + \epsilon \cdot \varphi) - F(p)$  by using  $f_\epsilon^*$  for both variational maximization objectives as follows,

$$\begin{aligned} & F(p + \epsilon \cdot \varphi) - F(p) \\ & \leq \left[ \int_{\mathcal{X}} f_\epsilon^*(x) \cdot [p(x) + \epsilon \cdot \varphi(x)] \, dx - F^*(f_\epsilon^*) \right] - \left[ \int_{\mathcal{X}} f_\epsilon^*(x) \cdot p(x) \, dx - F^*(f_\epsilon^*) \right] \\ & = \epsilon \cdot \int_{\mathcal{X}} f_\epsilon^*(x) \cdot \varphi(x) \, dx. \end{aligned} \quad (46)$$

Combining (45) and (46) with the sandwich theorem of limits [60], we obtain that  $F(p + \epsilon \cdot \varphi) - F(p)$  tends to zero as  $\epsilon$  goes to zero. Nevertheless, to derive a characterization for the right hand side of (43), it remains to quantify some distance measure between  $f_p^*$  and  $f_\epsilon^*$ . Fortunately, since  $F^*$  is strongly convex, we are able to set a constant  $\gamma > 0$  such that, for any two measurable functions  $f_1$  and  $f_2$ , we have

$$\int_{\mathcal{X}} \left[ \frac{\partial F^*}{\partial f_1}(x) - \frac{\partial F^*}{\partial f_2}(x) \right] \cdot [f_1(x) - f_2(x)] \, dx \geq \gamma \cdot \int_{\mathcal{X}} |f_1(x) - f_2(x)|^2 \, dx. \quad (47)$$

Moreover, since  $f_p^*$  and  $f_\epsilon^*$  are maximizers of the optimization problems in (4) and (44), respectively, we observe that  $\partial F^* / \partial f_p^* = p$  and  $\partial F^* / \partial f_\epsilon^* = p + \epsilon \cdot \varphi$ . Hence, by applying Cauchy-Schwarz inequality to the left hand side of (47), we have

$$\gamma \cdot \int_{\mathcal{X}} (f_\epsilon^* - f_p^*)^2 \, dx \leq \epsilon \cdot \int_{\mathcal{X}} \varphi \cdot (f_\epsilon^* - f_p^*) \, dx \leq \epsilon \cdot \left( \int_{\mathcal{X}} \varphi^2 \, dx \right)^{1/2} \cdot \left[ \int_{\mathcal{X}} (f_\epsilon^* - f_p^*)^2 \, dx \right]^{1/2}, \quad (48)$$

which implies that  $\|f_\epsilon^* - f_p^*\|_{\ell_2} \leq \epsilon \cdot \|\varphi\|_{\ell_2}$  and consequently  $f_\epsilon^*$  converges to  $f_p^*$  as  $\epsilon$  tends to zero. Then, the sandwich theorem of limits can be applied to the whole right hand side of (43), by plugging (45) and (46) into (43), to obtain that

$$\int_{\mathcal{X}} \frac{\delta F}{\delta p}(x) \cdot \varphi(x) \, dx = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \cdot [F(p + \epsilon \cdot \varphi) - F(p)] = \int_{\mathcal{X}} f_p^*(x) \cdot \varphi(x) \, dx$$

holds for any  $\varphi \in \mathcal{F}$ , which implies the result  $\delta F / \delta p = f_p^*$ .

To conclude, by combining (12) for  $F$  defined in (4), we obtain the final explicit form of the Riemannian gradient  $\text{grad } F = -\text{div}[p \cdot \nabla(f_p^*)]$ . Therefore, the proof of this proposition is completed.  $\blacksquare$

## C.2. Proof of Proposition 6

**Proof** According to the definition of exponential maps, to prove (13), it is sufficient to show the following results for a curve  $\gamma: [0, 1/h) \rightarrow \mathcal{M}$  defined by setting  $\gamma(t) = [\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_{\#} \mu$  for all  $t \in [0, 1/h)$ , where  $u \in \mathcal{F}$  and  $h$  denotes the Lipschitz constant of  $\nabla u$ .

- (i)  $\gamma(0) = \mu$ .

(ii)  $\gamma'(0) = s$ .

(iii)  $\gamma(t)$  is a geodesic on  $\mathcal{M}$  for  $t \in [0, 1/h)$ .

Please note that  $\gamma(0) = \text{id}_\# \mu = \mu$ , where  $\text{id}: \mathcal{X} \rightarrow \mathcal{X}$  is the identity mapping. First of all, we will adopt the following lemma to demonstrate that  $\gamma(t)$  is a geodesic on  $\mathcal{M}$ .

**Lemma 20** *Suppose that  $\mathcal{X}$  denotes  $\mathbb{R}^d$  or a closed convex subset of  $\mathbb{R}^d$  with periodic boundary conditions. Let  $u: \mathcal{X} \rightarrow \mathcal{X}$  be a twice continuously differentiable function over  $\mathcal{X}$  with a  $h$ -Lipschitz continuous gradient  $\nabla u: \mathcal{X} \rightarrow \mathcal{X}$ . Then, for any  $\mu \in \mathcal{M}$ , a curve  $\gamma: [0, 1/h) \rightarrow \mathcal{M}$  defined by  $\gamma(t) = [\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_\# \mu$  is a geodesic on  $\mathcal{M}$ .*

**Proof** We consider separately the two cases where (i)  $\mathcal{X}$  is  $\mathbb{R}^d$  or (ii)  $\mathcal{X}$  denotes a subset of  $\mathbb{R}^d$  with periodic boundary condition. The former case is first considered, where  $\gamma(t)$  can be formulated as (52).

(i)  $\mathcal{X}$  is  $\mathbb{R}^d$ . For completeness, we first need to verify that  $\gamma$  according to the definition is indeed a curve in  $\mathcal{M}$ , i.e., the pushforward maps admit  $\gamma(t) \in \mathcal{M}$  for all  $t \in [0, 1/h)$ . To proceed in this direction, we define a potential function

$$\varphi_t(x) = \|x\|_2^2/2 + t \cdot u(x) \quad (49)$$

for all  $t \in [0, 1/h)$ . As a result, we have the corresponding vector field  $\nabla \varphi_t = \text{id} + t \cdot \nabla u$  exactly induced by our pushforward mapping in the definition, which indicates that  $\gamma(t) = [\nabla \varphi_t]_\# \mu$ . Since  $\nabla u$  is  $h$ -Lipschitz continuous, we observe that  $\varphi_t$  is strongly convex for all  $t \in [0, 1/h)$ . In addition,  $\varphi_t$  is also twice continuously differentiable as  $u$  possesses continuous second-order derivative, which implies the Jacobian of  $\nabla \varphi_t$ , i.e.,  $\nabla^2 \varphi_t$  is continuous and positive definite. Therefore, we conclude that  $\nabla \varphi_t: \mathbb{R}^d \rightarrow \mathbb{R}^d$  is an invertible mapping. Thus,  $[\nabla \varphi_t]_\# \mu$  still lies in the distribution family which admits absolute continuity with respect to the Lebesgue measure and positive density everywhere. To summarize, it has been shown that  $\gamma(t) \in \mathcal{M}$  for all  $t \in [0, 1/h)$ .

It follows to prove that the curve  $\gamma$  on  $\mathcal{M}$  is a geodesic. To show this, we apply the Brenier's Theorem (see, for example, Theorem 2.12 in [63]) to conclude that there exists a unique optimal transport plan between  $\mu$  and  $\gamma(t)$ , which can be written as the gradient of a convex function  $\varphi$ . Meanwhile, the theorem further implies that  $\nabla \varphi$  serves as the unique gradient of some convex function such that  $\gamma(t)$  can be expressed as  $[\nabla \varphi]_\# p$ . Combining with the definition that  $\gamma(t) = [\nabla \varphi_t]_\# \mu$  for all  $t \in [0, 1/h)$ , we assert that  $\nabla \varphi_t$  represents the optimal transportation plan between  $p$  and  $\gamma(t)$ . With such useful characterization, we fix any  $\bar{t} \in [0, 1/h)$  below and show that  $\gamma$  is a geodesic when confined to  $[0, \bar{t}]$ . It is observed that

$$\nabla \varphi_t = \text{id} + t \cdot \nabla u = [1 - (t/\bar{t})] \cdot \text{id} + (t/\bar{t}) \cdot \nabla \varphi_{\bar{t}},$$

holds for any  $t \in [0, \bar{t}]$ , where  $\varphi_{\bar{t}}$  is a strongly convex function. Hence, we can write  $\gamma(t)$  in terms of  $\{[1 - (t/\bar{t})] \cdot \text{id} + (t/\bar{t}) \cdot \nabla \varphi_{\bar{t}}\}_\# \mu$ . To compute the Wasserstein distance of two points on the curve, for any  $0 \leq t_1 < t_2 \leq \bar{t}$ , we have

$$\begin{aligned} & W_2[\gamma(t_1), \gamma(t_2)] \\ &= \left[ \int_{\mathcal{X}} \left\| \{[1 - (t_1/\bar{t})]x + (t_1/\bar{t}) \cdot \nabla \varphi_{\bar{t}}(x)\} - \{[1 - (t_2/\bar{t})]x + (t_2/\bar{t}) \cdot \nabla \varphi_{\bar{t}}(x)\} \right\|^2 dp(x) \right]^{1/2} \\ &= (t_2 - t_1)/\bar{t} \cdot \left[ \int_{\mathbb{R}^d} \|x - \nabla \varphi_{\bar{t}}(x)\|^2 dp(x) \right]^{1/2} \\ &= (t_2 - t_1)/\bar{t} \cdot W_2[\mu, \gamma(\bar{t})], \end{aligned} \quad (50)$$



which implies that  $\{\gamma(t)\}_{t \in [0, \bar{t}]}$  is a reparametrized geodesic. Since  $\bar{t}$  is arbitrarily chosen within  $[0, 1/h)$ , it then holds that  $\gamma(t)$  is a geodesic for  $0 \leq t < 1/h$ .

(ii)  $\mathcal{X}$  denotes a subset of  $\mathbb{R}^d$  with periodic boundary condition. We are left to prove the lemma for the case where  $\mathcal{X}$  is a closed convex subset of  $\mathbb{R}^d$  with periodic boundary condition. In this case, any  $x \in \mathcal{X}$  can be identified with an equivalence class of  $\mathbb{R}^d$ . Moreover, each probability measure  $\mu \in \mathcal{P}(\mathcal{X})$  is unique identified with a periodic measure  $\bar{\mu} \in \mathcal{P}(\mathbb{R}^d)$  such that  $\bar{\mu}$  coincides with  $\mu$  on  $\mathcal{X}$ . We say  $\bar{\mu}$  is the periodic extension of  $\mu$ . Since  $\mu$  is absolutely continuous with respect to the Lebesgue measure and admits positive density, so is  $\bar{\mu}$ . Moreover,  $u: \mathcal{X} \rightarrow \mathbb{R}$  can also be extended as a periodic function on  $\mathbb{R}^d$ , and  $\varphi_t(x) = \|x\|_2^2/2 + t \cdot u(x)$  is a strongly convex, twice continuously differentiable, and periodic function on  $\mathbb{R}^d$ . Then it can be shown that  $(\text{id} + t \cdot \nabla u)_{\#} \bar{\mu}$  is the periodic extension of  $[\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_{\#} \mu$  [7]. Thus, these two measures coincide on  $\mathcal{X}$ , i.e.,

$$[\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_{\#} \mu = (\text{id} + t \cdot \nabla u)_{\#} \bar{\mu}|_{\mathcal{X}}, \quad (51)$$

where  $\cdot|_{\mathcal{X}}$  denotes the restriction to  $\mathcal{X}$ . Note that we have shown that  $[\nabla \phi_t]_{\#} \bar{\mu}$  is absolutely continuous with respect to the Lebesgue measure and has positive density. Thus, by restricting  $[\nabla \phi_t]_{\#} \bar{\mu}$  to  $\mathcal{X}$ , (51) implies that  $\gamma(t) \in \mathcal{M}$  for all  $t \in [0, 1/h)$ , i.e.,  $\gamma$  is a curve on  $\mathcal{M}$ .

Furthermore, to show that  $\gamma$  is a geodesic, we utilize the generalization of Brenier's theorem to probability distributions over a Riemannian manifold [25]. For any  $t \in [0, 1/h)$ , since  $\gamma(t) \in \mathcal{M}$ , there exists a unique optimal transportation plan  $\Upsilon: \mathcal{X} \rightarrow \mathcal{X}$  between  $\mu$  and  $\gamma(t)$  such that  $\gamma(t) = \Upsilon_{\#} \mu$ . Moreover,  $\Upsilon$  takes the form of  $\Upsilon(x) = \text{Exp}_x[-\nabla \psi(x)]$  for some  $\psi: \mathcal{X} \rightarrow \mathbb{R}$  such that  $\|x\|_2^2/2 - \psi(x)$  is convex. Hence, due to the uniqueness and the fact that  $\nabla u$  is  $h$ -Lipschitz,  $\text{Exp}_{\mathcal{X}}(-t \cdot \nabla u)$  is the optimal transportation plan between  $\mu$  and  $\gamma(t)$ .

Similarly in what follows we fix any  $\bar{t} \in [0, 1/H)$  and show that  $\gamma$  is a geodesic for  $t \in [0, \bar{t}]$ . For any  $0 \leq t_1 < t_2 \leq \bar{t}$ , following the derivations in (50) and combining (51), we have

$$\begin{aligned} W_2[\gamma(t_1), \gamma(t_2)] &= W_2\{[\text{Exp}_{\mathcal{X}}(t_1 \cdot \nabla u)]_{\#} \mu, [\text{Exp}_{\mathcal{X}}(t_2 \cdot \nabla u)]_{\#} \mu\} \\ &= W_2\left\{(\text{id} + t_1 \cdot \nabla u)_{\#} \bar{\mu}|_{\mathcal{X}}, (\text{id} + t_2 \cdot \nabla u)_{\#} \bar{\mu}|_{\mathcal{X}}\right\} = (t_2 - t_1)/\bar{t} \cdot W_2\left\{\bar{\mu}|_{\mathcal{X}}, (\text{id} + t_2 \cdot \nabla u)_{\#} \bar{\mu}|_{\mathcal{X}}\right\} \\ &= (t_2 - t_1)/\bar{t} \cdot W_2[\mu, \gamma(\bar{t})], \end{aligned}$$

where the second and the last equality follows from (51). Thus, we obtain that  $\{\gamma(t)\}_{t \in [0, \bar{t}]}$  is a geodesic up to reparametrization, which concludes the proof of Lemma 20.  $\blacksquare$

With the required Lemma in place, to finish the proof of Proposition 6, it remains to show that  $\gamma'(0) = s$ . Similar to the proof of the above lemma, we distinguish the two cases where  $\mathcal{X}$  is  $\mathbb{R}^d$  and  $\mathcal{X}$  is a closed convex subset of  $\mathbb{R}^d$  with periodic boundary condition. In the former case,

$$[\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_{\#} \mu = (\text{id} + t \cdot \nabla u)_{\#} \mu, \quad \forall t \in [0, 1/h). \quad (52)$$

To simplify the notation, we denote  $T_t = \text{id} + t \cdot \nabla u$ , which is invertible for  $t \in [0, 1/h)$ . By the definition of pushforward measures, we have

$$\gamma(t)(x) = [(T_t)_{\#} \mu](x) = \mu[T_t^{-1}(x)] \cdot \left| \frac{d}{dx} [T_t^{-1}(x)] \right|,$$

where the second equality follows from the change-of-variable formula and  $\left| \frac{d}{dx} [T_t^{-1}(x)] \right|$  is the determinant of the Jacobian. Moreover, when  $t$  is sufficiently small, for any  $x \in \mathcal{X}$ , Taylor expansion in  $t$  yields that

$T_t^{-1}(x) = x - t \cdot (\nabla u)(x) + o(t)$ , which further implies that

$$\begin{aligned}\gamma(t)(x) &= \mu[x - t \cdot (\nabla u)(x) + o(t)] \cdot |I_d - t \cdot (\nabla^2 u)(x) + o(t)| \\ &= [\mu(x) - t \cdot \langle (\nabla u)(x), (\nabla \mu)(x) \rangle + o(t)] \cdot \{1 - t \cdot \text{Tr}[(\nabla^2 u)(x)] + o(t)\} \\ &= \mu(x) - t \cdot \langle (\nabla u)(x), (\nabla \mu)(x) \rangle - t \cdot \text{Tr}[(\nabla^2 u)(x)] \cdot \mu(x) + o(t).\end{aligned}\quad (53)$$

where in the second equality we apply Taylor expansion to  $\mu[x - t \cdot (\nabla u)(x) + o(t)]$ . Moreover, since

$$\langle (\nabla u)(x), (\nabla p)(x) \rangle + \text{Trace}[(\nabla^2 u)(x)] \cdot p(x) = \text{div}[p(x) \cdot (\nabla u)(x)],$$

by (53) we obtain that

$$\gamma(t)(x) = \mu(x) - t \cdot \text{div}[p(x) \cdot (\nabla u)(x)] + o(t),$$

which implies that  $\gamma'(0) = -\text{div}(p \cdot \nabla u) = s$ .

It remains to show  $\gamma'(0) = -\text{div}(p \cdot \nabla u) = s$  when  $\mathcal{X}$  is a closed compact subset of  $\mathbb{R}^d$  with periodic boundary condition. As shown in the proof of Lemma 20,  $p$  can be periodically extended to a measure  $\bar{\mu}$  on  $\mathbb{R}^n$  and that such an extension is unique. Furthermore, the solution  $u$  to the elliptic equation  $-\text{div}(-p \cdot \nabla u) = s$  can also be viewed as a periodic function on  $\mathbb{R}^d$ . Then it can be shown that  $(\text{id} + t \cdot \nabla u)_\# \bar{\mu}$  is the periodic extension of  $[\text{Exp}_{\mathcal{X}}(t \cdot \nabla u)]_\# \mu$  and that (51) holds. Note that we have shown that  $\tilde{\gamma}(t) = (\text{id} + t \cdot \nabla u)_\# \bar{\mu}$  satisfies that  $\tilde{\gamma}(0) = \bar{\mu}$  and  $\tilde{\gamma}'(0) = s$ . Therefore, restricting to  $\mathcal{X}$ , we conclude that  $\gamma'(t) = s$ , which completes the proof of the proposition.  $\blacksquare$

### C.3. Examples for functionals with the variational form

The following example shows that if the entropy functional is  $f$ -divergence, the conjugate function  $F^*$  will be strongly smooth when the link function  $\psi$  is strongly convex, with respect to  $\ell_2$ -norm.

**Example 2** *Let  $p, q$  be two density function over a compact domain  $\mathcal{X}$ , then the  $f$ -divergence*

$$I_\psi(p, q) = \int_{\mathcal{X}} p(x) \cdot \psi\left(\frac{p(x)}{q(x)}\right) dx \quad (54)$$

with a strongly convex and smooth function  $\psi$  admits the following variational forms (as functionals of  $p$  and  $q$  respectively),

$$I_\psi(p, q) = I_{\psi, q}(p) = \sup_{f \in \mathcal{F}_p} \left\{ \int_{\mathcal{X}} f(x) p(x) dx - F_q^*(f) \right\}, \quad (55)$$

$$I_\psi(p, q) = I_{\psi, p}(q) = \sup_{f \in \mathcal{F}_q} \left\{ \int_{\mathcal{X}} f(x) q(x) dx - F_p^*(f) \right\}, \quad (56)$$

where

$$F_q^*(f) = \int_{\mathcal{X}} -(\psi^*)^{-1}(-f(x)) q(x) dx \quad (57)$$

and

$$F_p^*(f) = \int_{\mathcal{X}} \psi^*(f(x)) p(x) dx \quad (58)$$

are strongly convex and smooth functionals,  $\mathcal{F}_p$  and  $\mathcal{F}_q$  are the same set of all measurable functions on  $\mathcal{X}$ . We denote by  $\gamma$  its strong convexity parameter and by  $L$  the smoothness parameter.

**Proof** Following [46], Fenchel convex duality [55] ensures that we have  $I_\psi(p, q)$  in the form of the conjugate of  $\psi$  as:

$$I_\psi(p, q) = \sup_g \left\{ \int g(x)q(x) dx - \int \psi^*(g(x))p(x) dx \right\}, \quad (59)$$

where  $\psi^*$  is a strongly convex and smooth function since the convexity and smoothness of  $\psi$ , and the supremum is taken over all measurable function  $g : \mathcal{X} \rightarrow \mathbb{R}$ . Hence by replacing  $g$  with  $f$  formally we obtain (56).

Moreover, by letting  $f(x) = -\psi^*(g(x))$ , we have

$$I_\psi(p, q) = \sup_f \left\{ \int f(x)p(x) dx - \int -(\psi^*)^{-1}(-f(x))q(x) dx \right\}. \quad (60)$$

By setting  $F^*(f) = \int -(\psi^*)^{-1}(-f(x))q(x) dx$ , which can be verified to be strongly convex and smooth w.r.t. function  $f$ , we show the desired result in (55).  $\blacksquare$

Please note that by convex duality and optimality condition in Lemma 1 of [46], the function class  $\mathcal{F}_p$  ( $\mathcal{F}_q$ ) over which the supremum is taken can be restricted to a smaller one  $\tilde{\mathcal{F}}$  as long as  $\tilde{\mathcal{F}}$  contains the differential  $\partial I_\psi(p, q)/\partial p$  ( $\partial I_\psi(p, q)/\partial q$ ). In this sense, Assumption 7 for a smaller function class  $\tilde{\mathcal{F}}$  is feasible since it is natural for the smooth (with continuous first-order gradient and Hessian) function  $\partial I_\psi(p, q)/\partial q$  over a compact domain  $\mathcal{X}$  to possess a closed, bounded and equicontinuous gradient mapping by Weierstrass theorem [11].

## Appendix D. Proof of Convergence of Variational Form Maximization

We first illustrate our reverse Poincaré inequality with more details. To this end, we restate Lemma 21 and prove with a compact domain  $\mathcal{X}$  as follows.

### D.1. Approximation function class and reverse Poincaré inequality

Recall that  $\tilde{\mathcal{F}}$  is the function class of NNs, defined in (15).

**Lemma 21** *We consider a function class  $\tilde{\mathcal{F}}$  such that  $\nabla \tilde{\mathcal{F}}$  is closed, bounded, and equicontinuous. More precisely,*

- (a) “ $\nabla \tilde{\mathcal{F}}$  is bounded” means that there exists a positive constant  $M < \infty$  such that  $\|\nabla f(x)\| \leq M$  for each  $x \in \mathcal{X}$  and each  $f \in \tilde{\mathcal{F}}$ .
- (b) “ $\nabla \tilde{\mathcal{F}}$  is equicontinuous” implies that for every  $\epsilon > 0$  there exists  $\delta > 0$  (which depends only on  $\epsilon$ ) such that for any  $x, y \in \mathcal{X}$  with the metric  $d(\cdot, \cdot)$ , if  $d(x, y) < \delta$ , then it follows that

$$\|\nabla f(x) - \nabla f(y)\| < \epsilon, \quad \forall f \in \tilde{\mathcal{F}}. \quad (61)$$

Then we have for every  $p \geq 1$ , there exists a constant  $\tilde{K}$  such that

$$\int_{\mathcal{X}} \|\nabla f(x)\|^p d\mu \leq \tilde{K} \int_{\mathcal{X}} |f(x)|^p d\mu, \quad (62)$$

for any  $f \in \tilde{\mathcal{F}}$ , where  $\mathcal{X}$  is a compact subset of a metric space,  $\mu$  is a nonnegative measure over  $\mathcal{X}$ .

**Proof of Lemma 21**

**Proof** With the notations of  $L_\mu^p(\mathcal{X})$  norm, (62) can be rewritten as

$$\|\nabla f\|_{L_\mu^p(\mathcal{X})} \leq \tilde{K} \|f\|_{L_\mu^p(\mathcal{X})}, \quad (63)$$

On the other hand, Poincaré inequality (See, e.g., Chapter 5 of [18]) claims that there exists some constant  $K'$  such that

$$\|f - \frac{1}{\mu(\mathcal{X})} \int_{\mathcal{X}} f d\mu\|_{L_\mu^p(\mathcal{X})} \leq K' \|\nabla f\|_{L_\mu^p(\mathcal{X})}, \quad (64)$$

We define an equivalence relation  $\sim$  on  $L_\mu^p(\mathcal{X})$  such that for any  $f, g \in L_\mu^p(\mathcal{X})$ , we have

$$f \sim g \quad \text{if and only if} \quad f - g = C \quad (65)$$

for some constant  $C$ . Then we denote by  $\tilde{L}_\mu^p(\mathcal{X}) = L_\mu^p(\mathcal{X}) / \sim$  the new function space consisting of equivalence classes  $\tilde{f}$ 's of locally summable functions  $f : \mathcal{X} \rightarrow \mathbb{R}$ . Consequently, we define the norm in  $\tilde{L}_\mu^p(\mathcal{X})$  as  $\|\tilde{f}\|_{\tilde{L}_\mu^p(\mathcal{X})} = \|f\|_{L_\mu^p(\mathcal{X})}$ , and a new gradient operator  $\tilde{\nabla} : \tilde{L}_\mu^p(\mathcal{X}) \rightarrow \tilde{L}_\mu^p(\mathcal{X})$  such that for any  $\tilde{f} \in \tilde{L}_\mu^p(\mathcal{X})$ ,  $\tilde{\nabla}$  acts as  $\tilde{\nabla} \tilde{f} = \nabla f$ . Moreover, the inverse of  $\tilde{\nabla}$  exists, denoted by  $\tilde{\nabla}^{-1}$ .

Therefore, (64) implies that the inverse gradient operator  $\tilde{\nabla}^{-1}$  is continuous. Our goal is reduced to showing that the gradient operator is continuous, *i.e.*, the preimage of open sets in  $\tilde{L}_\mu^p(\mathcal{X})$  under such mapping are also open in  $\tilde{L}_\mu^p(\mathcal{X})$ .

According to (62) and the properties of continuous mappings between topological spaces (See, e.g., Appendix of [35]), for every compact subset  $X \subseteq \tilde{L}_\mu^p(\mathcal{X})$ , the image (preimage of  $\tilde{\nabla}$ ) under  $\tilde{\nabla}^{-1}$ , *i.e.*,  $\tilde{\nabla}^{-1}(X)$ , is compact in  $\tilde{L}_\mu^p(\mathcal{X})$ , and closed in  $\tilde{L}_\mu^p(\mathcal{X})$  since  $\tilde{L}_\mu^p(\mathcal{X})$  is a Hausdorff space. On the other hand, by Arzela-Ascoli Theorem the set  $\tilde{L}_\mu^p(\mathcal{X}) \cap \nabla \tilde{\mathcal{F}}$  is compact. Then for every closed subset  $Y \subseteq \tilde{L}_\mu^p(\mathcal{X}) \cap \nabla \tilde{\mathcal{F}}$ , we have  $Y$  is also a compact subset, of which the image under  $\tilde{\nabla}^{-1}$  is closed in  $\tilde{L}_\mu^p(\mathcal{X})$ . According to the definition the continuity of  $\tilde{\nabla}$  is shown, hence (62) is proved. ■

**Remark 22** *Specifically, we remark that the conditions above which enable the gradient bound (Lemma 21) to hold keep consistent with all the other restrictions on function class  $\tilde{\mathcal{F}}$ , and can be easily realized by a general class of neural networks.*

*First, the small function class  $\tilde{\mathcal{F}}$  is designed to solve VFM defined in (4) numerically efficiently and meanwhile to provide a decent characterization of Riemannian gradient estimation error bounds. We also observe that to achieve an equivalence between pushing particles and the exponential map, functions in this smaller class are also required to admit uniformly  $h$ -Lipschitz continuous gradients. Fortunately, the equicontinuity of  $\nabla \tilde{\mathcal{F}}$  can be implied by  $h$ -Lipschitzness of  $\nabla f \in \nabla \tilde{\mathcal{F}}$ . In other words, the overall assumptions of function class  $\tilde{\mathcal{F}}$  for our whole analysis are concluded as below.*

**Assumption 23 (Overall assumptions of approximation function class)** *The class of functions  $\tilde{\mathcal{F}}$  over a compact domain  $\mathcal{X}$  satisfies the condition that  $\nabla \tilde{\mathcal{F}}$  is closed, bounded, and for each  $f \in \tilde{\mathcal{F}}$ ,  $\nabla f$  is  $h$ -Lipschitz continuous with  $h > 0$ .*

*Then we check that our neural network parametrized function class is indeed qualified for such  $\tilde{\mathcal{F}}$ . Recall that without loss of generality, we parametrize a function  $f : \mathcal{X} \rightarrow \mathbb{R}$ , *i.e.*, the decision variable for VFM, by the class of two-layer neural networks below, which is denoted as  $\text{NN}(\beta; w)$ , with each member function as*

$$f_\beta(x) = \frac{1}{\sqrt{w}} \sum_{i=1}^w b_i \cdot \sigma([\beta]_i^\top x), \quad (66)$$

where  $x \in \mathcal{X}$  denotes the input data point,  $w$  gives the width of the neural network,  $b_i \in \{-1, 1\}$  for  $i \in [w]$  denotes the output weights,  $\sigma(\cdot)$  is a smooth activation function, and  $\beta = ([\beta]_1^\top, \dots, [\beta]_w^\top)^\top \in \mathbb{R}^{wd}$  with  $[\beta]_i \in \mathbb{R}^d$  ( $i \in [w]$ ) are the overall input weights. For initialization, we consider the random strategy

$$b_i \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(\{-1, 1\}), \quad [\beta(0)]_i \stackrel{\text{i.i.d.}}{\sim} N(0, I_d/(wd)), \quad \text{for all } i \in [w]. \quad (67)$$

Note that for technical analysis reason, we restrict the input weights  $\beta$  to an bounded  $\ell_2$ -ball centered at the initializer  $\beta(0)$  by an additional projection step  $\Pi_{\mathcal{B}^0(r_f)}(\bar{\beta}) = \operatorname{argmin}_{\beta \in \mathcal{B}^0(r_f)} \{\|\beta - \bar{\beta}\|_2\}$ , where  $\mathcal{B}^0(r_f) = \{\beta : \|\beta - \beta(0)\|_2 \leq r_f\}$ . During the training process, we only backpropagate w.r.t.  $\beta$ , while keeping  $b_i$  ( $i \in [w]$ ) intact at the initialization which accounts for the feasibility to omit the dependency on  $b_i$  ( $i \in [w]$ ) in  $\text{NN}(\beta; w)$  and  $f_\beta(x)$  in what follows. Therefore, we can directly observe that each  $f_\beta \in \text{NN}(\beta; w)$  is closed, and the  $\ell_2$ -norm of the gradient  $\nabla_\beta f_\beta$  is always bounded over a compact domain  $\mathcal{X}$ . Furthermore, we require the Hessian for  $f_\beta$  w.r.t. parameter  $\beta \in \mathbb{R}^{wd}$  to possess a bounded norm, which is easy to check for neural networks. It turns out that the qualified function class encompasses a wide range of normal neural networks without “sharp points” or “high frequency oscillation” as functions.

## D.2. Statistical error of approximating the solution of VFM

In what follows, we proceed to derive the gradient error bound. According to (62) of Lemma 21, we turn to bound the approximation error of a two-layer neural network for the following variational form maximization (VFM) problem defined in (14),

$$\max_{\beta \in \mathbb{R}^{wd}} \left\{ \frac{1}{N} \sum_{i=1}^N f_\beta(x_i) - F^*(f_\beta) \right\}. \quad (68)$$

Note that for notational simplicity and applicability of results to both players, in this section we omit the references to  $\mu$  and  $\nu$ . To emphasize we fixed the  $b_i$ 's throughout the training, we write  $f_\beta(x)$  while omitting the dependency on  $b_i$ 's. We first show that the overparameterization of the NN,  $f_\beta$  parametrized by  $\beta$ , guarantees that it behaves as its local linearization at the random initialization  $\beta(0)$ . To this end, we define

$$f_\beta^0(x) = \frac{1}{\sqrt{w}} \sum_{i=1}^w b_i \cdot \sigma'([\beta(0)]_i^\top x) \cdot [\beta]_i^\top x, \quad (69)$$

whereby the linear structure of  $f_\beta^0(x)$  indicates

$$f_\beta^0(x) = \langle \nabla_\beta f_\beta^0(x), \beta \rangle. \quad (70)$$

We write  $\beta(s)$  as the value of the parameter  $\beta$  at the  $s$ -th iteration of VFM (Algorithm 3). For the simplicity of notations, we denote by

$$G_{\beta(s)}(x) = -\nabla_\beta f_{\beta(s)}(x) + \nabla_\beta F^*(f_{\beta(s)}) \quad (71)$$

the stochastic gradient vector. Similarly, we also define

$$G_{\beta(s)}^0(x) = -\nabla_\beta f_{\beta(s)}^0(x) + \nabla_\beta F^*(f_{\beta(s)}^0). \quad (72)$$

Furthermore, let  $\mu$  be the probability measure corresponding to input data distribution for the NNs. We denote by

$$\bar{G}_{\beta(s)} = \mathbb{E}_\mu [G_{\beta(s)}(x)] = \int_{\mathcal{X}} G_{\beta(s)}(x) d\mu(x) \quad (73)$$

the population mean of the stochastic gradient vector at the  $s$ -th iteration, and by  $\bar{G}_{\beta(s)}^0$  its localization version. Without loss of generality, we assume  $\|x\| \leq 1$  for the data.

We first present the following assumption for bounding the variance of the stochastic gradient vector  $G_{\beta(s)}(x)$ , which casts the analysis of global convergence into tracking the mean of the gradient vector. We denote by  $\mathbb{E}_{\text{init}}[\cdot]$  the expectation over the random initialization for neural network parameter  $\beta$ , and  $\mathbb{E}_\mu[\cdot]$  the expectation over the input  $x \in \mathcal{X}$  conditioned on the random initialization.

**Assumption 24 (Variance of the Stochastic Update Vector)** *For any  $s \leq t$ , there exists a constant  $C_G^2 = \mathcal{O}(r_f^2)$  independent of  $s$  such that*

$$\mathbb{E}_{\text{init}} \|G_{\beta(s)} - \bar{G}_{\beta(s)}\|_{L_\mu^2(\mathcal{X})}^2 \leq C_G^2. \quad (74)$$

The following theorem provides the final characterization of the gradient error. Recall that in Algorithm 3,  $t$  is the number of iterations as well as the sample size denoted by  $N$  for either  $N_\mu$  or  $N_\nu$ ,  $w$  is the width of neural networks, and  $r_f$  is the projection radius.

Furthermore, to formalize the smoothness of the activation function, we have the following assumption.

**Assumption 25 (Smooth Activation Function)** *There exists an absolute constant  $h > 0$  such that for any  $x_1, x_2 \in \mathbb{R}^d$  we have*

$$|\sigma'(x_1) - \sigma'(x_2)| \leq h \|x_1 - x_2\|. \quad (75)$$

For technical consideration, we also assume  $\sigma(0) = 0$ .

The following lemma quantifies the variance by the introduced function  $f_\beta^0$  and the generic function  $f_\beta$  in terms of  $r_f$  and  $w$ .

**Lemma 26** *Under Assumption 24 and 25, given  $\beta \in \mathcal{B}^0(r_f)$ , we have*

$$\mathbb{E}_{\text{init}} \|f_\beta - f_\beta^0\|_{L_\mu^2(\mathcal{X})}^2 = \mathcal{O}(w^{-1}r_f^4). \quad (76)$$

**Proof** By the definition of  $f_\beta$  and  $f_\beta^0$ , we have

$$\begin{aligned} & |f_\beta(x) - f_\beta^0(x)| \\ & \leq \frac{1}{\sqrt{w}} \left| \sum_{i=1}^w (\sigma([\beta]_i^\top x) - \sigma'([\beta(0)]_i^\top x) [\beta]_i^\top x) \right| \\ & \leq \frac{1}{\sqrt{w}} \sum_{i=1}^w \left| \sigma([\beta]_i^\top x) - \sigma'([\beta(0)]_i^\top x) [\beta]_i^\top x \right| \\ & \leq \frac{1}{\sqrt{w}} \sum_{i=1}^w \left| \sigma([\beta(0)]_i^\top x) - \sigma'([\beta(0)]_i^\top x) [\beta(0)]_i^\top x \right| \\ & \quad + \frac{1}{\sqrt{w}} \sum_{i=1}^w \left| \left( \int_0^1 (\sigma'((1-\eta)[\beta(0)]_i^\top x + \eta[\beta]_i^\top x)) \cdot d\eta \right) \cdot x^\top ([\beta]_i - [\beta(0)]_i) \right|, \end{aligned} \quad (77)$$

where the last inequality holds by the Taylor expansion, the second last inequality follows from the triangle inequality, and the second equality holds as  $|b_i| = 1$ . The first term on the right-hand side of (77) is bounded by

$$\begin{aligned} \frac{1}{\sqrt{w}} \sum_{i=1}^w \left| \sigma([\beta(0)]_i^\top x) - \sigma'([\beta(0)]_i^\top x) [\beta(0)]_i^\top x \right| & \leq \mathcal{O} \left( \frac{1}{\sqrt{w}} \mathbb{E}_{\text{init}} \sum_{i=1}^w \|[\beta(0)]_i\|^2 \right) \\ & = \mathcal{O} \left( \frac{1}{\sqrt{w}} \right), \end{aligned} \quad (78)$$

where we use the Taylor expansion of  $\sigma$  at 0 and the fact that  $\sigma(0) = 0$ .

By squaring both sides of (77) and applying Assumption 25 to the right-hand side of (77), we obtain

$$\begin{aligned}
\mathbb{E}_{\text{init}, \mu} |f_\beta(x) - f_\beta^0(x)|^2 &= \mathcal{O}\left(\frac{1}{w}\right) + \mathcal{O}\left(\left(\frac{h}{\sqrt{w}} \mathbb{E}_{\text{init}} \sum_{i=1}^w \|[\beta]_i - [\beta(0)]_i\|^2\right)^2\right) \\
&= \mathcal{O}\left(\left(\frac{h}{\sqrt{w}} \sum_{i=1}^w \|[\beta]_i - [\beta(0)]_i\|^2\right)^2\right) \\
&= \mathcal{O}\left(\frac{h}{w} \|\beta - \beta(0)\|^4\right), \\
&= \mathcal{O}\left(\frac{r_f^4}{w}\right). \tag{79}
\end{aligned}$$

Hence, it follows that

$$\mathbb{E}_{\text{init}} \|f_\beta - f_\beta^0\|_{L_\mu^2(\mathcal{X})}^2 = \mathcal{O}(w^{-1}r_f^4), \tag{80}$$

which concludes the proof.  $\blacksquare$

The following lemma characterizes the difference between the expected gradients of the the original neural network approximator and the locally linearized one.

**Lemma 27** *For any  $0 \leq s \leq t$ , we have the following linear approximation error at each iteration:*

$$\mathbb{E}_{\text{init}} \|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2^2 = \mathcal{O}(w^{-1}r_f^4 + w^{-1}r_f^2). \tag{81}$$

**Proof** By the definition of  $\bar{G}_{\beta(s)}$  and  $\bar{G}_{\beta(s)}^0$ , we have

$$\begin{aligned}
&\mathbb{E}_{\text{init}} \|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2^2 \\
&\leq \mathbb{E}_{\text{init}, \mu} \|\underbrace{-\nabla_\beta f_{\beta(s)}(x) + \nabla_\beta F^*(f_{\beta(s)}) + \nabla_\beta f_{\beta(s)}^0(x) - \nabla_\beta F^*(f_{\beta(s)}^0)}_{\text{(i)}}\|_2^2 \\
&\leq 2 \underbrace{\mathbb{E}_{\text{init}, \mu} \|\nabla_\beta f_{\beta(s)}(x) - \nabla_\beta f_{\beta(s)}^0(x)\|_2^2}_{\text{(i)}} + 2 \underbrace{\mathbb{E}_{\text{init}, \mu} \|\nabla_\beta F^*(f_{\beta(s)}) - \nabla_\beta F^*(f_{\beta(s)}^0)\|_2^2}_{\text{(ii)}}. \tag{82}
\end{aligned}$$

In what follows, we upper bound term (i) and (ii) respectively.

**Upper Bounding (i):** Recall that

$$\nabla_\beta f_\beta(x) = 1/\sqrt{w} \cdot (b_1 \cdot \sigma'([\beta]_1^\top x) \cdot x^\top, \dots, b_w \cdot \sigma'([\beta]_w^\top x) \cdot x^\top)^\top,$$

and

$$\nabla_\beta f_\beta^0(x) = 1/\sqrt{w} \cdot (b_1 \cdot \sigma'([\beta(0)]_1^\top x) \cdot x^\top, \dots, b_w \cdot \sigma'([\beta(0)]_w^\top x) \cdot x^\top)^\top.$$

We have

$$\begin{aligned}
\|\nabla_\beta f_{\beta(s)}(x) - \nabla_\beta f_{\beta(s)}^0(x)\|_2^2 &= \frac{1}{w} \sum_{i=1}^w (\sigma'([\beta]_i^\top x) - \sigma'([\beta(0)]_i^\top x))^2 \cdot \|x\|^2 \\
&\leq \frac{h^2}{w} \sum_{i=1}^w \|[\beta]_i - [\beta(0)]_i\|^2 \\
&= \frac{h^2 r_f^2}{w}. \tag{83}
\end{aligned}$$

Here the last inequality follows from the condition  $\|x\|^2 \leq 1$ , which is also used to derive (77). By taking expectation on (83) we obtain

$$\mathbb{E}_{\text{init}, \mu} \|\nabla_{\beta} f_{\beta(s)}(x) - \nabla_{\beta} f_{\beta(s)}^0(x)\|_2^2 = \mathcal{O}(w^{-1} r_f^2). \quad (84)$$

Therefore, we have bounded term (i) in (82).

**Upper Bounding (ii):** By the  $L$ -smoothness of the dual functional  $F^*$ , it follows that

$$\begin{aligned} & \mathbb{E}_{\text{init}, \mu} \|\nabla_{\beta} F^*(f_{\beta(s)}) - \nabla_{\beta} F^*(f_{\beta(s)}^0)\|_2^2 \\ & \leq \mathbb{E}_{\text{init}, \mu} \|f_{\beta(s)}(x) - f_{\beta(s)}^0(x)\|_2^2, \end{aligned} \quad (85)$$

where the right-hand side of (85) is exactly the left-hand side of (76). Hence by Lemma 26, the term (ii) is bounded by  $\mathcal{O}(w^{-1} r_f^4)$ . Combining (i) and (ii) concludes the proof for Lemma 27.  $\blacksquare$

In what follows, with the explicit linearized learning target  $f_{\beta^*}^0$ , we are able to characterize the global convergence of Algorithm 3 by the difference between output estimated function and  $f_{\beta^*}^0$ .

**Lemma 28** *Suppose that  $t > 64$  iterations of Algorithm 1 are run, and the stepsize is set to be  $\eta = t^{-1/2}$ . Then, under Assumption 24 and 25 we have*

$$\mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2 = \mathcal{O}(r_f^2 t^{-1/2} + w^{-1/2} r_f^3 + w^{-1} r_f^4), \quad (86)$$

where  $\beta^*$  is the approximate stationary point such that

$$\beta^* = \Pi_{\mathcal{B}^0(r_f)}(\beta^* - \eta \bar{G}_{\beta^*}^0), \quad (87)$$

and  $\hat{\beta} = 1/t \cdot \sum_{s=0}^{t-1} \beta(s)$ .

**Proof** First we bound the progress of the one-step update. By the convexity of  $\mathcal{B}^0(r_f)$  and the approximate stationary condition (87), for each  $s < t$  we have

$$\begin{aligned} & \mathbb{E}_{\mu} [\|\beta(s+1) - \beta^*\|_2^2 | \beta(s)] \\ & = \mathbb{E}_{\mu} \left[ \left\| \Pi_{\mathcal{B}^0(r_f)}(\beta(s) - \eta G_{\beta(s)}) - \Pi_{\mathcal{B}^0(r_f)}(\beta^* - \eta \bar{G}_{\beta^*}^0) \right\|_2^2 | \beta(s) \right] \\ & \leq \mathbb{E}_{\mu} \left[ \left\| (\beta(s) - \beta^*) - \eta (G_{\beta(s)} - \bar{G}_{\beta^*}^0) \right\|_2^2 | \beta(s) \right] \\ & = \|\beta(s) - \beta^*\|_2^2 - 2\eta \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)} - \bar{G}_{\beta^*}^0 \rangle + \eta^2 \|G_{\beta(s)} - \bar{G}_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2. \end{aligned} \quad (88)$$

Then our target is reduced to upper bound the last two terms above. For the inner product term, by applying Hölder's inequality we have

$$\begin{aligned} \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)} - \bar{G}_{\beta^*}^0 \rangle & = \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0 \rangle + \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0 \rangle \\ & \geq -\|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2 \cdot \|\beta(s) - \beta^*\|_2 + \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0 \rangle \\ & \stackrel{(a)}{\geq} -r_f \|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2 + \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0 \rangle, \end{aligned} \quad (89)$$

where (a) uses the fact that  $\|\beta(s) - \beta^*\|_2 \leq r_f$ . For the rest inner product term in (89), by plugging in the definition of stochastic gradient vectors we obtain

$$\begin{aligned} \langle \beta(s) - \beta^*, \bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0 \rangle & = \mathbb{E}_{\mu} \left\langle \beta(s) - \beta^*, \nabla_{\beta} f_{\beta(s)}^0(x) - \nabla_{\beta} f_{\beta(s)}^0(x) \right\rangle \\ & \quad + \mathbb{E}_{\mu} \left\langle \beta(s) - \beta^*, \nabla_{\beta} F^*(f_{\beta(s)}^0) - \nabla_{\beta} F^*(f_{\beta^*}^0) \right\rangle \\ & = \mathbb{E}_{\mu} \left\langle \beta(s) - \beta^*, \nabla_{\beta} F^*(f_{\beta(s)}^0) - \nabla_{\beta} F^*(f_{\beta^*}^0) \right\rangle, \end{aligned} \quad (90)$$



where we use the fact that  $\nabla_{\beta} f_{\beta}^0(x)$  is independent of the value of  $\beta$ . We proceed to derive the lower bound of (??) by the explicit functional gradient form:

$$\begin{aligned}
& \mathbb{E}_{\mu} \left\langle \nabla_{\beta} F \left( f_{\beta(s)}^0 \right) - \nabla_{\beta} F \left( f_{\beta^*}^0 \right), \beta(s) - \beta^* \right\rangle \\
&= \mathbb{E}_{\mu} \left\langle \left( \frac{\delta F^*}{\delta f} \left( f_{\beta(s)}^0 \right) - \frac{\delta F^*}{\delta f} \left( f_{\beta^*}^0 \right) \right) \cdot \nabla_{\beta} f_{\beta}^0, \beta(s) - \beta^* \right\rangle \\
&= \mathbb{E}_{\mu} \left\langle \frac{\delta F^*}{\delta f} \left( f_{\beta(s)}^0 \right) - \frac{\delta F^*}{\delta f} \left( f_{\beta^*}^0 \right), f_{\beta(s)}^0 - f_{\beta^*}^0 \right\rangle \\
&\stackrel{(b)}{\geq} \gamma \|f_{\beta(s)}^0 - f_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2,
\end{aligned} \tag{91}$$

where (b) is due to the  $\gamma$ -strong convexity of functional  $F^*$ .

On the other hand, the third norm term in (88) is estimated using the Cauchy-Schwarz inequality as follows,

$$\begin{aligned}
\|G_{\beta(s)} - \bar{G}_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2 &\leq 2\|G_{\beta(s)} - \bar{G}_{\beta(s)}\|_{L_{\mu}^2(\mathcal{X})}^2 + 2\|\bar{G}_{\beta(s)} - \bar{G}_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2 \\
&\leq 2\mathbb{E}_{\text{init}}\|G_{\beta(s)} - \bar{G}_{\beta(s)}\|_{L_{\mu}^2(\mathcal{X})}^2 + 4\|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2^2 + 4\|\bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0\|_2^2
\end{aligned} \tag{92}$$

where the first term herein is upper bounded by the variance of  $G_{\beta(s)}$  in (74) of Assumption 24, and the second one is controlled by lemma 26. Thus it suffices to upper bound the last term, *i.e.*, the squared difference between the expected gradient of time-step  $s$  and the optimal one of linearization approximation. We have

$$\begin{aligned}
\|\bar{G}_{\beta(s)}^0 - \bar{G}_{\beta^*}^0\|_2^2 &= \|\nabla_{\beta} \mathbb{E} f_{\beta^*}^0 - \nabla_{\beta} \mathbb{E} f_{\beta(s)}^0 + \nabla_{\beta} F^*(f_{\beta(s)}^0) - \nabla_{\beta} F^*(f_{\beta^*}^0)\|_2^2 \\
&\leq 2\|\nabla_{\beta} \mathbb{E} f_{\beta^*}^0 - \nabla_{\beta} \mathbb{E} f_{\beta(s)}^0\|_2^2 + 2\|\nabla_{\beta} F^*(f_{\beta(s)}^0) - \nabla_{\beta} F^*(f_{\beta^*}^0)\|_2^2 \\
&\leq 2\|f_{\beta^*}^0 - f_{\beta(s)}^0\|_{L_{\mu}^2(\mathcal{X})}^2 + 2L\|f_{\beta^*}^0 - f_{\beta(s)}^0\|_{L_{\mu}^2(\mathcal{X})}^2
\end{aligned} \tag{93}$$

$$= 2(1+L)\|f_{\beta^*}^0 - f_{\beta(s)}^0\|_{L_{\mu}^2(\mathcal{X})}^2, \tag{94}$$

where (93) follows from the  $L$ -smoothness of the entropy dual functional by Example 2. To conclude, we have

$$\begin{aligned}
& \mathbb{E}_{\mu} [\|\beta(s+1) - \beta^*\|_2^2 \mid \beta(s)] \\
&\leq \|\beta(s) - \beta^*\|_2^2 + 2\eta r_f \|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2 \\
&\quad + 2\eta^2 \mathbb{E}_{\text{init}} \|G_{\beta(s)} - \bar{G}_{\beta(s)}\|_{L_{\mu}^2(\mathcal{X})}^2 + 4\eta^2 \|\bar{G}_{\beta(s)} - \bar{G}_{\beta(s)}^0\|_2^2 \\
&\quad + 2\eta(4\eta(1+L) - \gamma) \|f_{\beta(s)}^0 - f_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2.
\end{aligned} \tag{95}$$

By rearranging (95), we have

$$\begin{aligned}
\|f_{\beta(s)} - f_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2 &\leq 2\|f_{\beta(s)} - f_{\beta(s)}^0\|_{L_{\mu}^2(\mathcal{X})}^2 + 2\|f_{\beta(s)}^0 - f_{\beta^*}^0\|_{L_{\mu}^2(\mathcal{X})}^2 \\
&\leq (\gamma\eta - 4\eta^2(L+1))^{-1} \left( \|\beta(s) - \beta^*\|_2^2 - \mathbb{E}_{\mu} [\|\beta(s+1) - \beta^*\|_2^2 \mid \beta(s)] \right. \\
&\quad \left. + 2C_G^2\eta^2 + \mathcal{O}\left(w^{-1/2}r_f^3 + w^{-1}r_f^4\right) \right),
\end{aligned} \tag{96}$$

where the second inequality follows from lemma 26 and 27, as well as the fact  $\eta < \gamma/8(L+1)$  resulted from  $t > 64(L+1)^2/\gamma^2$  and  $\eta = t^{-1/2}$ . We proceed to take total expectation on both sides of (96) and

telescoping for  $s + 1 \in [t]$  ( $t \geq 1$ ) to obtain

$$\begin{aligned}
\mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}^0\|_{L_\mu^2(\mathcal{X})}^2 &= \mathbb{E}_{\text{init}, \mu} [(f_{\hat{\beta}}(x) - f_{\beta^*}^0(x))^2] \\
&\leq \frac{1}{t} \sum_{s=0}^{t-1} \mathbb{E}_{\text{init}, \mu} [(f_{\beta(s)}(x) - f_{\beta^*}^0(x))^2] \\
&\leq t^{-1} \cdot (\gamma\eta - 4\eta^2(L+1))^{-1} \cdot \left( \mathbb{E}_{\text{init}} [\|\beta(0) - \beta^*\|_2^2] + 2tC_G^2\eta^2 \right. \\
&\quad \left. + \mathcal{O}(w^{-1/2}r_f^3 + w^{-1}r_f^4) \right). \tag{97}
\end{aligned}$$

By plugging the conditions on  $t$  and  $\eta$ , we have

$$\begin{aligned}
t^{-1} \cdot (\gamma\eta - 4\eta^2(L+1))^{-1} &= t^{-1/2} \cdot \frac{1}{\gamma - 4\eta(L+1)} \\
&\stackrel{\eta < \gamma/8(L+1)}{\leq} t^{-1/2} \cdot \frac{1}{\gamma - \gamma/2} \\
&= \frac{2}{\gamma\sqrt{t}}, \tag{98}
\end{aligned}$$

Then, we obtain the following bound,

$$\begin{aligned}
\mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}^0\|_{L_\mu^2(\mathcal{X})}^2 &\leq \frac{2}{\gamma\sqrt{t}} (\mathbb{E}_{\text{init}} [\|\beta(0) - \beta^*\|_2^2] + 2C_G^2) + \mathcal{O}(r_f^{5/2}w^{-1/4} + r_f^3w^{-1/2}) \tag{99}
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{2(r_f^2 + 2C_G^2)}{\gamma\sqrt{t}} + \mathcal{O}(r_f^{5/2}w^{-1/4} + r_f^3w^{-1/2}) \\
&= \mathcal{O}(r_f^2t^{-1/2} + w^{-1/2}r_f^3 + w^{-1}r_f^4). \tag{100}
\end{aligned}$$

Hence we complete the proof.  $\blacksquare$

Now it is straightforward to prove the main theorem for VFM with neural network parametrized function class as follows.

### D.3. Gradient Error of Neural Networks

**Theorem 29** *Under Assumption 7, 24 and 25, within the  $k$ -th iteration of Algorithm 2, the gradient error  $\bar{\varepsilon}_k$  defined in (20) satisfies*

$$\bar{\varepsilon}_k = \bar{\varepsilon}_k(N) = \mathcal{O}(\bar{\varepsilon}(N)) = \mathcal{O}\left(\tilde{K}\left(\frac{r_f^2}{N^{1/2}} + \frac{r_f^3}{w^{1/2}} + \frac{r_f^4}{w}\right)\right). \tag{101}$$

We remark that the order of the gradient error is independent of iteration  $k$  and can be decomposed into the generalization error of  $\mathcal{O}(r_f^2/N^{1/2})$  for SGD over the neural tangent kernel and the error of  $\mathcal{O}(r_f^3/w^{1/2})$ ,  $\mathcal{O}(r_f^4/w)$  for approximating the neural network by a linear function. In particular, the overall gradient errors decay to zero at the rate of  $1/\sqrt{N}$  with a sufficiently large width  $w$  of the neural network.

**Proof** By Jensen's inequality with respect to squared  $L_\mu^2$ -norm, we have

$$\begin{aligned} & \mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}\|_{L_\mu^2(\mathcal{X})}^2 \\ & \leq 2\mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}^0\|_{L_\mu^2(\mathcal{X})}^2 + 2\mathbb{E}_{\text{init}} \|f_{\beta^*}^0 - f_{\beta^*}\|_{L_\mu^2(\mathcal{X})}^2. \end{aligned} \quad (102)$$

On one hand, by Lemma 28 the first term above can be bounded with  $\mathcal{O}(r_f^2 t^{-1/2} + w^{-1/2} r_f^3 + w^{-1} r_f^4)$ . On the other hand, the second term in (102) is bounded by  $\mathcal{O}(r_f^4 w^{-1})$  through setting  $\beta = \beta^*$  in Lemma 26. To conclude, the total bound for (102) turns out to be  $\mathcal{O}(r_f^2 t^{-1/2} + w^{-1/2} r_f^3 + w^{-1} r_f^4)$ , which completes the proof for the first part of the theorem.

To verify (101), we invoke Lemma 21 by setting  $f$  to  $\tilde{f}_k^* - \hat{f}_k^*$  with all possible iteration  $k$  for both players and  $p$  to 2, combining with (20) to obtain for any  $k \in \mathbb{N}$ , we have

$$\begin{aligned} \bar{\epsilon}_k &= \mathbb{E}_{X_0} \int_{\mathcal{X}} \|\nabla \tilde{f}_k^*(x) - \nabla \hat{f}_k^*(x)\|_2^2 d\tilde{\rho}_k \\ &\stackrel{(62)}{\leq} \tilde{K} \mathbb{E}_{X_0} \int_{\mathcal{X}} |\tilde{f}_k^*(x) - \hat{f}_k^*(x)|_2^2 d\tilde{\rho}_k \\ &= \tilde{K} \mathbb{E}_{X_0} \mathbb{E}_{\text{init}} \|f_{\hat{\beta}} - f_{\beta^*}\|_{L_{\tilde{\rho}_k}^2(\mathcal{X})}^2 \\ &= \mathcal{O}\left(\tilde{K} \left(\frac{r_f^2}{N^{1/2}} + \frac{r_f^3}{w^{1/2}} + \frac{r_f^4}{w}\right)\right), \end{aligned} \quad (103)$$

where the last equality follows from (102) and the measure  $\tilde{\rho}_k$  accounts for either distribution iterates at some iteration  $k$ ,  $\tilde{\mu}_k$  or  $\tilde{\nu}_k$  in the paper. Therefore, we conclude the whole proof of the theorem for the statistical errors of VFM steps.  $\blacksquare$

## Appendix E. Proof of Convergence for Distributional Game Optimization

In this section, we layout the complete proof needed to reach the convergence rate of particle-based infinite-dimensional game optimization, for which some intermediate lemmas are also listed below. We define  $\nu^*(\mu) \in \mathcal{I}_F(\mu) \triangleq \operatorname{argmax}_\nu F(\mu, \nu)$  for simplicity. Also note that according to the definition in Algorithm 1, we have  $\nu_{k+1} = \nu_{K_\nu}(\mu_k)$  as the last iterate of the inner loop for the  $k$ -th outer loop. Moreover, in the formal description of VTIG in Algorithm 1, we adopt sets of discrete particles to represent the underlying distribution iterates being updated, where  $X_l^\nu(\tilde{\mu}_k)$  denotes the set of particles for player  $\nu$  at the  $l$ -th iteration given  $\tilde{\mu}_k$  as the current distribution iterate for player  $\mu$ . Similarly for the formal proof, we will use the distribution notation  $\nu_l(\tilde{\mu}_k)$  indicating the distribution iterate derived from constructed transportation maps  $\{T_{k,l}\}_{k \in [K_\mu], l \in [K_\nu]}$  given  $\tilde{\mu}_k$ .

### E.1. Proof of Proposition 30

Considering the interaction of the two players, our algorithm runs multiple gradient ascent steps in the inner loop to estimate the *inner maximization value functional* defined as  $G(\mu) \triangleq \max_{\nu \in \mathcal{M}(\mathcal{X}_\nu)} F(\mu, \nu)$ , of which the Riemannian gradient *w.r.t.*  $\mu$  at the optimum  $\nu^*(\mu)$  is adopted to estimate the Riemannian gradient of  $G(\mu)$ . Inspired by this, we rewrite (3) as  $\min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)$  and observe that the overall VTIG behaves as a gradient descent-like algorithm over the inner maximization value functional  $G$ . Hence we can make use of nonconvex optimization techniques to analyze the overall convergence properties of such zero-sum two-player games. For simplicity we define  $H_\mu(\nu) \triangleq -F(\mu, \nu)$ . The following proposition identifies that  $\mathcal{M}(\mathcal{X})$  is well defined.

**Proposition 30** *The manifold  $\mathcal{M}(\mathcal{X})$  is compact, that is, there exists  $R > 0$  such that for any  $\mu_1, \mu_2 \in \mathcal{M}(\mathcal{X})$  we have  $W_2(\mu_1, \mu_2) \leq 2R$ .*

**Proof** Since  $\mathcal{X}$  is defined as a compact domain, according to the topological properties in Theorem 6.16 of [64], the manifold  $\mathcal{M}(\mathcal{X})$  is also compact. Also, as the underlying variable space  $\mathcal{X}$  is bounded, the Wasserstein distance defined over  $\mathcal{M}(\mathcal{X})$  over  $\mathcal{X}$  is bounded, by setting  $2R = \sup_{\mu_1, \mu_2 \in \mathcal{M}(\mathcal{X}_\mu)} \mathcal{W}_2(\mu_1, \mu_2)$  we arrive at the conclusion. ■

Before proceeding, we first state the following additional definitions and lemmas.

**Definition 31** *We say that a function  $G : \mathcal{M}(\mathcal{X}) \rightarrow \mathbb{R}$  satisfies the Quadratic Growth (QG) condition w.r.t.  $\mathcal{W}_2$  metric with constant  $\gamma > 0$  if*

$$G(\mu) - G(\mu^*) \geq \frac{\gamma}{2} W_2^2(\mu, \mu^*), \quad \forall \mu \in \mathcal{M}(\mathcal{X}), \quad (104)$$

where  $\mu^*$  is the minimizer of the function. Note that if function  $G$  is PL with constant  $\xi$ , then  $G$  satisfies the QG condition with constant  $\gamma = 4\xi$ .

The next lemma demonstrates the stability of  $\nu^*(\mu)$  controlled by the variation of  $\mu$ .

**Lemma 32** *Given that  $\mathcal{I}_F(\mu)$  is closed, then for any  $\mu_1, \mu_2$  and  $\nu_1 \in \mathcal{I}_F(\mu_1)$ , there exists a  $\nu_2 \in \mathcal{I}_F(\mu_2)$  such that*

$$W_2(\nu_1, \nu_2) \leq \frac{L_0}{2\xi} W_2(\mu_1, \mu_2). \quad (105)$$

**Proof** By the Lipschitzness condition 2 and the PL condition of  $F$ , we have

$$G(\mu_2) - F(\mu_2, \nu_1) \leq \frac{L_0^2}{2\xi} W_2^2(\mu_1, \mu_2). \quad (106)$$

Moreover, by the QG property of  $H_\mu(\nu)$  it follows that there exists a  $\nu_2 \in \mathcal{I}_F(\mu_2)$  such that

$$W_2(\nu_1, \nu_2) \leq \frac{L_0}{2\xi} W_2(\mu_1, \mu_2), \quad (107)$$

which concludes the result. ■

At the two-player game optimization scale (VFM not involved), Considering the interaction of the two players, our algorithm runs multiple gradient ascent steps in the inner loop to estimate the *inner maximization value functional* defined as  $G(\mu) \triangleq \max_{\nu \in \mathcal{M}(\mathcal{X}_\nu)} F(\mu, \nu)$ , of which the Riemannian gradient w.r.t.  $\mu$  at the optimum  $\nu^*(\mu)$  is adopted to estimate the Riemannian gradient of  $G(\mu)$ . Inspired by this, we rewrite (3) as  $\min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)$  and observe that the overall VTIG behaves as a gradient descent-like algorithm over the inner maximization value functional  $G$ . Hence we can make use of nonconvex optimization techniques to analyze the overall convergence properties of such min-max two-player games. For simplicity we define  $H_\mu(\nu) \triangleq -F(\mu, \nu)$ . The following proposition identifies that  $\mathcal{M}(\mathcal{X})$  is well defined.

It is not straightforward to see  $\text{grad } G(\mu) = \text{grad}_\mu F(\mu, \nu^*(\mu))$  as there may be multiple  $\nu^*(\mu)$ 's under Assumption 3, breaking the condition of Danskin's theorem [5] even for finite cases. Fortunately, we show in the following with Riemannian PL condition, we can still prove a Danskin-type result as the generalization of the original theorem. In a nutshell, our proposed algorithm is a gradient descent-like algorithm on the inner maximization value function. To proceed with two-player game of in a higher hierarchy for Problem (3), we follow the standard assumptions of smoothness in different aspect of  $F$  and provide a Danskin-type lemma in the following section.

## E.2. Danskin-type Lemma

**Lemma 33** Under Assumption 2 and 3, we have

$$\text{grad } G(\mu) = \text{grad}_\mu F(\mu, \nu^*(\mu)), \quad \nu^*(\mu) \in \underset{\nu \in \mathcal{M}(\mathcal{X}_\nu)}{\text{argmin}} H_\mu(\nu).$$

In addition,  $G$  is  $\tilde{L}_1$ -Lipschitz and  $\tilde{L}$ -smooth with  $\tilde{L}_1 = L_\mu + L_\nu L_0 / (2\xi)$ ,  $\tilde{L} = L_1 + L_0^2 / (2\xi)$ .

**Proof** By the definition of the directional derivative and gradient in Section B.2, we take the curve, which starts with  $\gamma(0) = \mu \in \mathcal{M}(\mathcal{X})$  and  $\gamma'(0) = u \in T_\mu \mathcal{M}(\mathcal{X})$ , as the geodesic. Then by Taylor series expansion at  $\gamma(0)$ , we have for any scalar  $\tau$ :

$$\begin{aligned} G[\gamma(\tau)] - G(\mu) &= F(\gamma(\tau), \nu^*(\gamma(\tau))) - F(\mu, \nu^*(\mu)) \\ &= \langle \text{grad}_\mu F(\mu, \nu^*(\mu)), \tau u \rangle_\mu + \left\langle \text{grad}_\nu F(\mu, \nu^*(\mu)), \text{Exp}_{\nu^*(\mu)}^{-1}(\nu^*(\gamma(\tau))) \right\rangle_{\nu^*(\mu)} + \mathcal{O}(\tau^2) \\ &= \langle \text{grad}_\mu F(\mu, \nu^*(\mu)), \tau u \rangle_\mu + \mathcal{O}(\tau^2). \end{aligned} \quad (108)$$

where the last equality follows from the optimal condition on  $\nu$ :  $\text{grad}_\nu F(\mu, \nu^*(\mu)) = 0$ , and the additional higher order term in  $\tau$  integrates terms in  $\mathcal{W}_2(\nu^*(\gamma(\tau)), \nu^*(\mu))$  since by (107) there exists  $\nu^*(\gamma(\tau)) \in \mathcal{I}_F(\gamma(\tau))$  such that

$$\mathcal{W}_2(\nu^*(\gamma(\tau)), \nu^*(\mu)) \leq \frac{L_0}{2\xi} \tau \|u\|. \quad (109)$$

On the other hand, by the limit form of directional derivative we have

$$\begin{aligned} \langle \text{grad } G(\mu), u \rangle_\mu &= \nabla_u G(\mu) \\ &= \frac{d}{d\tau} G[\gamma(\tau)] \Big|_{t=0} \\ &= \lim_{\tau \rightarrow 0^+} \frac{G[\gamma(\tau)] - G[\gamma(0)]}{\tau} \\ &= \langle \text{grad}_\mu F(\mu, \nu^*(\mu)), \tau u \rangle_\mu. \end{aligned} \quad (110)$$

Due to the arbitrariness of  $u$ , we conclude that  $\text{grad } G(\mu) = \text{grad}_\mu F(\mu, \nu^*(\mu))$ .

In what follows we show the Lipschitzness and smoothness of  $G$ . For  $\mu_1, \mu_2 \in \mathcal{M}(\mathcal{X})$ , let  $\nu^*(\mu_1) \in \mathcal{I}_F(\mu_1)$  and  $\nu^*(\mu_2) \in \underset{\nu \in \mathcal{I}_F(\mu_2)}{\text{argmin}} \mathcal{W}_2^2(\nu, \nu^*(\mu_1))$ , then

$$\begin{aligned} &|G(\mu_1) - G(\mu_2)| \\ &= |F(\mu_1, \nu^*(\mu_1)) - F(\mu_2, \nu^*(\mu_1)) + F(\mu_2, \nu^*(\mu_1)) - F(\mu_2, \nu^*(\mu_2))| \\ &\leq L_\mu \mathcal{W}_2(\mu_1, \mu_2) + L_\nu \mathcal{W}_2(\nu^*(\mu_1), \nu^*(\mu_2)) \\ &\leq \left( L_\mu + \frac{L_\nu L_0}{2\xi} \right) \mathcal{W}_2(\mu_1, \mu_2), \end{aligned} \quad (111)$$

where the last inequality holds by (107). Hence by setting  $\tilde{L}_1 = L_\mu + L_\nu L_0/(2\xi)$  we obtain that function  $G$  is  $\tilde{L}_1$ -Lipschitz. In addition, for the difference between gradients we have

$$\begin{aligned}
\mathbf{d}(\text{grad } G(\mu_1), \text{grad } G(\mu_2)) &= \mathbf{d}(\text{grad}_\mu F(\mu_1, \nu^*(\mu_1)), \text{grad}_\mu F(\mu_2, \nu^*(\mu_2))) \\
&\leq \mathbf{d}(\text{grad}_\mu F(\mu_1, \nu^*(\mu_1)), \text{grad}_\mu F(\mu_2, \nu^*(\mu_1))) \\
&\quad + \mathbf{d}(\text{grad}_\mu F(\mu_2, \nu^*(\mu_1)), \text{grad}_\mu F(\mu_2, \nu^*(\mu_2))) \\
&\leq \tilde{L}_1 \mathcal{W}_2(\mu_1, \mu_2) + L_0 \mathcal{W}_2(\nu^*(\mu_1), \nu^*(\mu_2)) \\
&\leq \left( \tilde{L}_1 + \frac{L_0^2}{2\xi} \right) \mathcal{W}_2(\mu_1, \mu_2), \tag{112}
\end{aligned}$$

where the second inequality holds by Assumption 2, and the last inequality still follows from (107). Therefore, by setting  $\tilde{L} = \tilde{L}_1 + L_0^2/(2\xi)$  we concludes the proof.  $\blacksquare$

### E.3. Convergence Results for Player $\nu$

Considering the nested loops in Algorithm 2, we first present the following linear convergence rate with statistical errors for player  $\nu$ . Recall that  $\tilde{\mathcal{F}}$  defined in (15) is the function class over which we solve (4). We define  $\nu^*(\mu) = \text{argmax}_{\nu \in \mathcal{M}} F(\mu, \nu)$  given  $\mu \in \mathcal{M}$ . Also, we write  $\tilde{\nu}_l(\tilde{\mu}_k)$  as  $\tilde{\nu}_l$  given any fixed  $k \geq 0$  for notational simplicity.

**Lemma 34** *Let  $F$  admit the variational form under Assumptions 2 and 3. Suppose that  $\tilde{\mathcal{F}}$  satisfies Assumption 7. Also, we set the stepsize  $\eta^\nu \in (0, \min\{1/(4L_\nu), 1/h\})$ . Then, for any  $l \geq 1$  and a fixed  $\mu \in \mathcal{M}$ , we have*

$$F(\mu, \nu^*(\mu)) - \mathbb{E}[F(\mu, \tilde{\nu}_l)] \leq \sigma^l \cdot [F(\mu, \nu^*(\mu)) - F(\mu, \tilde{\nu}_0)] + \eta^\nu \sigma^l \sum_{m=0}^{l-1} \bar{\varepsilon}_m \sigma^{-(m+1)}, \tag{113}$$

where  $\sigma = 1 - \xi\eta^\nu/2 \in (0, 1)$  is the contraction coefficient, the expectation is taken w.r.t. the initial samples  $X_0^\nu = \{x_{i,0}^\nu\}_{i \in [N_\nu]}$ , and  $\bar{\varepsilon}_m$  ( $m \in [l-1]$ ) is the (expected) gradient error at timestep  $m$  defined in (20).

**Proof** Our proof is based on quantifying some contraction between errors of adjacent iterates. Recall that we denote  $H_\mu(\nu) = -F(\mu, \nu)$ , and for fixed  $\mu \in \mathcal{M}(\mathcal{X})$ , We write  $H_\mu(\nu)$  as  $H(\nu)$  below. To begin with, by Proposition 6, we can equivalently write the iteration of VTIG for  $l \geq 0$  and  $\tilde{\nu}_l$  as

$$\tilde{\nu}_{l+1} = \text{Exp}_{\tilde{\nu}_l} \left\{ -\eta^\nu \cdot [\text{grad } H(\tilde{\nu}_l) + \delta_l] \right\}, \quad \delta_l = -\text{div}[\tilde{\nu}_l \cdot (\nabla \tilde{f}_l^* - \nabla f_l^*)]. \tag{114}$$

Notice that  $\delta_l \in T_{\tilde{\nu}_l} \mathcal{M}$  is a tangent vector at point  $\tilde{\nu}_l$ . Moreover, by Assumption 2 on  $F$  function  $H$  is  $L_2$ -smooth. Combining this property with (114), we have

$$\begin{aligned}
H(\tilde{\nu}_{l+1}) &\leq H(\tilde{\nu}_l) - \eta^\nu \cdot \langle \text{grad } H(\tilde{\nu}_l), \text{grad } H(\tilde{\nu}_l) \rangle_{\tilde{\nu}_l} - \eta^\nu \langle \text{grad } H(\tilde{\nu}_l), \delta_l \rangle_{\tilde{\nu}_l} \\
&\quad + \frac{(\eta^\nu)^2 L_2}{2} \langle \text{grad } H(\tilde{\nu}_l) + \delta_l, \text{grad } H(\tilde{\nu}_l) + \delta_l \rangle_{\tilde{\nu}_l} \\
&= H(\tilde{\nu}_l) - \left( \eta^\nu - \frac{(\eta^\nu)^2 L_2}{2} \right) \cdot \langle \text{grad } H(\tilde{\nu}_l), \text{grad } H(\tilde{\nu}_l) \rangle_{\tilde{\nu}_l} \\
&\quad + (\eta^\nu + (\eta^\nu)^2 \cdot L_2) \cdot |\langle \text{grad } H(\tilde{\nu}_l), \delta_l \rangle_{\tilde{\nu}_l}| + \frac{(\eta^\nu)^2 L_2}{2} \langle \delta_l, \delta_l \rangle_{\tilde{\nu}_l}, \tag{115}
\end{aligned}$$

where  $\langle \cdot, \cdot \rangle_{\tilde{\nu}_l}$  is the Riemannian metric of  $\mathcal{M}$  at  $\tilde{\nu}_l$ . By basic inequality  $2ab \leq a^2 + b^2$ , we have

$$|\langle \text{grad } H(\tilde{\nu}_l), \delta_l \rangle_{\tilde{\nu}_l}| \leq \frac{1}{2} \langle \text{grad } H(\tilde{\nu}_l), \text{grad } H(\tilde{\nu}_l) \rangle_{\tilde{\nu}_l} + \frac{1}{2} \langle \delta_l, \delta_l \rangle_{\tilde{\nu}_l}. \tag{116}$$

Thus, plugging (116) into (115), we obtain

$$H(\tilde{\nu}_{l+1}) \leq H(\tilde{\nu}_l) - \frac{\eta^\nu(1 - 2\eta^\nu L_2)}{2} \cdot \langle \text{grad } H(\tilde{\nu}_l), \text{grad } H(\tilde{\nu}_l) \rangle_{\tilde{\nu}_l} + \frac{\eta^\nu(1 + 2\eta^\nu L_2)}{2} \cdot \langle \delta_l, \delta_l \rangle_{\tilde{\nu}_l}. \quad (117)$$

Furthermore, since  $H$  is  $\xi$ -gradient dominated under Assumption 3, based on (117) we have

$$H(\tilde{\nu}_{l+1}) \leq H(\tilde{\nu}_l) - \xi\eta^\nu \cdot (1 - 2\eta^\nu L_2) \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] + \frac{\eta^\nu(1 + 2\eta^\nu L_2)}{2} \cdot \langle \delta_l, \delta_l \rangle_{\tilde{\nu}_l}. \quad (118)$$

As we have introduced in Section 2.1 and 4.1,  $\langle \delta_l, \delta_l \rangle_{\tilde{\nu}_l}$  is equal to  $\varepsilon_l$  defined in (20). Thus, (118) can be equivalently written as

$$H(\tilde{\nu}_{l+1}) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu) \leq [1 - \xi\eta^\nu \cdot (1 - 2\eta^\nu L_2)] \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] + \frac{\eta^\nu(1 + 2\eta^\nu L_2)}{2} \cdot \varepsilon_l. \quad (119)$$

Hence, we have derived the performance of a single step of variational transport with regard to the objective value.

Moreover, recall that we set the stepsize for player  $\nu$  to be a constant  $\eta^\nu \leq 1/(4L_2)$ , which guarantees

$$1 - \xi \cdot \eta^\nu \cdot (1 - 2\eta^\nu L_2) \leq 1 - \eta^\nu \cdot \xi/2 \in (0, 1), \quad (1 + 2\eta^\nu L_2)/2 \leq 1. \quad (120)$$

For simplicity of the notation, we define  $\sigma = 1 - \eta^\nu \cdot \xi/2$ . Thus, by (120), it follows that

$$H(\tilde{\nu}_{l+1}) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu) \leq \sigma \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] + \eta^\nu \varepsilon_l. \quad (121)$$

By multiplying  $\sigma^{-(l+1)}$  to both sides of (121), we obtain

$$\sigma^{-(l+1)} \cdot [H(\tilde{\nu}_{l+1}) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] \leq \sigma^{-l} \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] + \sigma^{-(l+1)} \cdot \eta^\nu \varepsilon_l. \quad (122)$$

Therefore,  $\{\sigma^{-l} \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)]\}_{l \geq 0}$  admits a telescoping sequence and thus by summing (122) over  $l$  it holds that

$$\sigma^{-l} \cdot [H(\tilde{\nu}_l) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)] \leq \sum_{m=0}^{l-1} \sigma^{-(m+1)} \cdot \eta^\nu \cdot \varepsilon_m + [H(\tilde{\nu}_0) - \inf_{\nu \in \mathcal{M}(\mathcal{X})} H(\nu)], \quad (123)$$

which is equivalent to

$$\sigma^{-l} \cdot [F(\mu, \tilde{\nu}^*(\mu)) - F(\mu, \tilde{\nu}_l)] \leq \sum_{m=0}^{l-1} \sigma^{-(m+1)} \cdot \eta^\nu \cdot \varepsilon_m + [F(\mu, \tilde{\nu}^*(\mu)) - F(\mu, \tilde{\nu}_0)] \quad (124)$$

for all  $l \geq 1$ , which implies the desired result by taking expectation with respect to the initial particle sampling  $X_0^\nu$  for player  $\nu$ .  $\blacksquare$

Lemma 34 characterizes the expected error of the inner loop by the sum of an optimization error decaying at a linear rate and a statistical error term of the order  $\mathcal{O}(N_\nu^{-1/2})$  according to §4.1. Moreover, (113) justifies that the particle-based functional gradient descent in  $W_2$ -space for a Riemannian PL objective behaves similarly as in finite-dimensional spaces [58] up to a scaled statistical error.

By Integrating the convergence result for one-player variational transport processes, together with the properties of relation between the inner loop and outer loop as well as objective landscape, we are now ready to demonstrate the main theorem for infinite-dimensional distributional game optimization.

#### E.4. Proof of Theorem 8

**Theorem 35 (Convergence of Infinite-Dimensional PL Games, Formal)** *Suppose that the objective  $F$  admits a variational form under Assumption 2 and 3. Also, the function class  $\mathcal{F}$  satisfies Assumption 7. We set the stepsizes to be  $\eta^\mu \in [0, \min\{1/h, 2/\widehat{L}\})$  and  $\eta^\nu \in (0, \min\{1/(4L_\nu), 1/h\})$ , where  $\widehat{L} = L_1 + L_0^2/\xi$ . Then, for any  $\theta > 0$ , if*

$$K_\nu \geq K_\nu(\theta) = \mathcal{O}\left(\log \frac{(1-\sigma)\widehat{M}_H - \eta^\nu \bar{\varepsilon}_\nu}{\theta} / \log \frac{1}{\sigma}\right), \quad \text{where } \widehat{M}_H = \max\left\{M_H, \frac{\eta^\nu \bar{\varepsilon}_\nu + 1}{1-\sigma}\right\}, \quad (125)$$

there exists an iteration  $k \in [K_\mu]$  such that

$$\mathbb{E}_{X_0}[\mathcal{J}_\mu^2(\tilde{\mu}_k, \tilde{\nu}_{k+1})] = \mathcal{O}\left(\frac{2}{\widehat{L}}(\Delta + \sqrt{\bar{\varepsilon}_\mu} + 2\tilde{L}R + L_G)^2 \cdot \left(2R(\Delta + \sqrt{\bar{\varepsilon}_\mu}) + \frac{M_G}{K_\mu}\right)\right), \quad (126)$$

$$\mathbb{E}_{X_0}[\mathcal{J}_\nu(\tilde{\mu}_k, \tilde{\nu}_{k+1})] = \mathcal{O}\left(\frac{L_2\Delta}{L_0}\right). \quad (127)$$

Here  $\Delta = L_0\sqrt{\frac{\eta^\nu \bar{\varepsilon}_\nu + \theta}{2\xi(1-\sigma)}}$ ,  $\widehat{L} = 1/\eta^\mu - \tilde{L}/2$ ,  $R = \sup_{\mu_1, \mu_2 \in \mathcal{M}} \mathcal{W}_2(\mu_1, \mu_2)/2$ , and the gradient error terms  $\bar{\varepsilon}_\mu$  and  $\bar{\varepsilon}_\nu$  are characterized in (21).

**Proof** [Proof of Theorem 8] We write  $\widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = -\text{div}(\tilde{\mu}_k \cdot \nabla \tilde{f}_k^*)$ , where  $\nabla \tilde{f}_k^*$  is the solution to the VFM problem in timestep  $k$ , as the gradient estimate w.r.t.  $\mu$  at timestep  $k$ . By the particle pushing step in Algorithm 2 we have

$$\langle \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} = -\frac{1}{\eta^\mu} \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k}. \quad (128)$$

It follows that

$$\begin{aligned} & \langle \text{grad}_\mu F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} \\ &= \langle \text{grad}_\mu F(\tilde{\mu}_k, \tilde{\nu}^*(\tilde{\mu}_k)) - \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} \\ & \quad - \frac{1}{\eta^\mu} \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} \\ &= \langle \Delta_k - \delta_k, \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} - \frac{1}{\eta^\mu} \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k}, \end{aligned} \quad (129)$$

where  $\nu^*(\tilde{\mu}_k) \in \mathcal{I}_F(\tilde{\mu}_k)$  and

$$\delta_k = -\text{div}[\tilde{\mu}_k \cdot (\nabla \tilde{f}_k^* - \nabla f_k^*)], \quad (130)$$

$$\Delta_k = \text{grad} G(\tilde{\mu}_k) - \text{grad}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}). \quad (131)$$

Hence, we have  $\Delta_k - \delta_k = \text{grad}_\mu F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)) - \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = \text{grad} G(\tilde{\mu}_k) - \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1})$  for any  $k \in [K_\mu]$ . Note that the error term  $\Delta_k$  is incurred by using  $\tilde{\nu}_{k+1}$  to approximate  $\nu^*(\tilde{\mu}_k)$ .

By Assumption 7,  $\nabla f$  is  $h$ -Lipschitz continuous on  $\mathcal{X}$  for all  $f \in \tilde{\mathcal{F}}$ . Since  $\eta^\mu < 1/h$ , by Proposition 6, we can equivalently write one timestep of the outer loop in VTIG as

$$\begin{aligned} \tilde{\mu}_{k+1} &= \text{Exp}_{\tilde{\mu}_k} \left\{ -\eta^\mu \cdot \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) \right\} \\ &= \text{Exp}_{\tilde{\mu}_k} \left\{ -\eta^\mu \cdot [\text{grad}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) + \delta_k] \right\} \end{aligned} \quad (132)$$

$$= \text{Exp}_{\tilde{\mu}_k} \left\{ -\eta^\mu \cdot [\text{grad} G(\tilde{\mu}_k) - \Delta_k + \delta_k] \right\}, \quad (133)$$



Note that  $\delta_k \in T_{\tilde{\mu}_k} \mathcal{M}$  is a tangent vector at point  $\tilde{\mu}_k$ . Since  $G$  is  $\tilde{L}$ -smooth under Lemma 33, combining (129) and (132), we obtain

$$G(\tilde{\mu}_{k+1}) \leq G(\tilde{\mu}_k) + \langle \text{grad}_{\mu} F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} + \frac{\tilde{L}}{2} \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} \quad (134)$$

$$\stackrel{(129)}{=} G(\tilde{\mu}_k) + \langle \Delta_k - \delta_k, \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} - \left( \frac{1}{\eta^\mu} - \frac{\tilde{L}}{2} \right) \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k}.$$

Since  $\eta^\mu < 2/\tilde{L}$ , a lower bound of  $\|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\|$  is needed. On the other hand, we have for any  $\mu \in \mathcal{M}$ ,

$$\begin{aligned} \langle \widehat{\text{grad}}_{\mu} F(\tilde{\mu}_k, \tilde{\nu}_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\mu) \rangle_{\tilde{\mu}_k} &\geq -\|\widehat{\text{grad}}_{\mu} F(\tilde{\mu}_k, \tilde{\nu}_{k+1})\|_{\tilde{\mu}_k} \cdot \|\text{Exp}_{\tilde{\mu}_k}^{-1}(\mu)\|_{\mu} \\ &\geq -\|\widehat{\text{grad}}_{\mu} F(\tilde{\mu}_k, \tilde{\nu}_{k+1})\|_{\tilde{\mu}_k} (\|\text{Exp}_{\tilde{\mu}_{k+1}}^{-1}(\mu)\|_{\mu} + \|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\|_{\tilde{\mu}_{k+1}}) \\ &\geq -\frac{1}{\eta^\mu} \|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\|_{\tilde{\mu}_{k+1}} \|\text{Exp}_{\tilde{\mu}_{k+1}}^{-1}(\mu)\|_{\mu} \\ &\quad - (\|\Delta_k\| + \|\delta_k\| + \|\text{grad } G(\tilde{\mu}_k)\|) \|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\|_{\tilde{\mu}_{k+1}}, \end{aligned} \quad (135)$$

where the first inequality follows from Cauchy-Schwartz inequality (135) is implied by triangle inequality *w.r.t.* Riemannian metric, and the last one follows from (128) and (129). Then by definitions  $\|\text{grad } G(\tilde{\mu}_k)\| \leq G_{\max}$  and  $\|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\| \leq 2R$ . Additionally, by Theorem 34 using fixed  $\mathbb{E}\epsilon_m = \bar{\epsilon}_\nu$  with expectation *w.r.t.* initial sampled particles, we have

$$\begin{aligned} \mathbb{E}\|\Delta_k\| &\leq L_0 \mathbb{E}W_2(\tilde{\nu}_{k+1}, \nu^*(\tilde{\mu}_k)) \\ &\leq L_0 \sqrt{\frac{\sigma^{K_\nu} [F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)) - F(\tilde{\mu}_k, \tilde{\nu}_0)] + \eta^\nu \bar{\epsilon}_\nu \cdot \frac{1-\sigma^{K_\nu}}{1-\sigma}}{2\xi}} \\ &\leq L_0 \sqrt{\frac{\eta^\nu \bar{\epsilon}_\nu + \theta}{2\xi(1-\sigma)}}, \end{aligned} \quad (136)$$

where the second inequality holds by applying Definition 31 to the gradient dominated function  $H_\mu(\nu) = -F(\mu, \nu)$  together with Theorem 34, and the last inequality follows from the choice of  $K_\nu$  in the theorem. Then by  $\|\text{grad } G(\tilde{\mu}_k)\| \leq G_{\max}$  and  $\|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\| \leq 2R$ , combining (135) we obtain

$$-\mathcal{J}_\mu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) \geq -(\|\Delta_k\| + \|\delta_k\| + G_{\max} + 2R/\eta^\mu) \|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\|, \quad (137)$$

that is,

$$\|\text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1})\| \geq \frac{\mathcal{J}_\mu(\tilde{\mu}_k, \tilde{\nu}_{k+1})}{\|\Delta_k\| + \|\delta_k\| + G_{\max} + 2\tilde{L}R}. \quad (138)$$

Plugging (138) into (134), we obtain the following progress made by pushing  $\mu$ -particles in the outer loop.

$$\begin{aligned} G(\mu_{k+1}) &\leq G(\tilde{\mu}_k) + \langle \Delta_k - \delta_k, \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} - \frac{(1/\eta^\mu - \tilde{L}/2) \mathcal{J}_\mu^2(\tilde{\mu}_k, \tilde{\nu}_{k+1})}{\left( \|\Delta_k\| + \|\delta_k\| + G_{\max} + 2\tilde{L}R \right)^2} \\ &\leq G(\tilde{\mu}_k) + 2R(\|\Delta_k\| + \|\delta_k\|) - \frac{(1/\eta^\mu - \tilde{L}/2) \mathcal{J}_\mu^2(\tilde{\mu}_k, \tilde{\nu}_{k+1})}{\left( \|\Delta_k\| + \|\delta_k\| + G_{\max} + 2\tilde{L}R \right)^2}, \end{aligned} \quad (139)$$

where the last inequality holds by applying the Cauchy-Schwartz inequality to the inner product term  $\langle \Delta_k - \delta_k, \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k}$  and  $\tilde{\mu}_k$ 's lie in a ball of radius  $R$ .

Therefore, we have a telescoping sequence and by summing (139) over  $k$  for the whole loop while taking expectation with respect to initial sampled particles on both sides we get

$$\begin{aligned} & \frac{1}{K_\mu} \sum_{k=0}^{K_\mu} \mathbb{E} \mathcal{J}_\mu^2(\tilde{\mu}_k, \tilde{\nu}_{k+1}) \\ & \leq \frac{2}{1/\eta^\mu - \tilde{L}/2} \mathbb{E} \left( 2\tilde{L} + G_{\max} + \|\Delta_k\| + \|\delta_k\| \right)^2 \cdot \left( 2R(\|\Delta_k\| + \|\delta_k\|) + \frac{G(\tilde{\mu}_0) - G(\tilde{\mu}_{K_\mu})}{K_\mu} \right) \\ & \leq \mathcal{O} \left( \frac{2}{1/\eta^\mu - \tilde{L}/2} \left( \Delta + \sqrt{\bar{\varepsilon}_\mu} + 2\tilde{L}R + G_{\max} \right)^2 \cdot \left( 2R(\Delta + \sqrt{\bar{\varepsilon}_\mu}) + \frac{M_G}{K_\mu} \right) \right), \end{aligned} \quad (140)$$

(141)

where the last inequality follows from (136). Here

$$\Delta = L_0 \sqrt{\frac{\eta_\nu \bar{\varepsilon}_\nu + \theta}{2\xi(1-\sigma)}}, \quad (142)$$

and

$$\sqrt{\bar{\varepsilon}_k} = \mathcal{O} \left( \sqrt{\tilde{K} \left( \frac{r_f^2}{N^{1/2}} + \frac{r_f^3}{w^{1/2}} + \frac{r_f^4}{w} \right)} \right).$$

Therefore, by the Pigeonhole principle there must exist  $k \in [K_\mu]$  such that

$$\mathbb{E} \mathcal{J}_\mu^2(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = \mathcal{O} \left( \frac{2}{1/\eta^\mu - \tilde{L}/2} \left( \Delta + \sqrt{\bar{\varepsilon}_\mu} + 2\tilde{L}R + G_{\max} \right)^2 \cdot \left( 2R(\Delta + \sqrt{\bar{\varepsilon}_\mu}) + \frac{M_G}{K_\mu} \right) \right).$$

Hence we complete the proof of the first term in (23) of Theorem 8.

For the second result on  $\mathcal{J}_\nu(\mu_k, \nu_{k+1})$ , we follow the similar technique by the Cauchy-Schwartz inequality and smoothness of function  $F$  with respect to  $\nu$  in Assumption 2 to obtain

$$\begin{aligned} \mathbb{E} \mathcal{J}_\nu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) & \leq \mathbb{E} \|\text{grad}_\nu F(\tilde{\mu}_k, \tilde{\nu}_{k+1})\|_{\tilde{\nu}_{k+1}} \\ & = \mathbb{E} \mathbf{d}(\text{grad}_\nu F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)), \text{grad}_\nu F(\tilde{\mu}_k, \tilde{\nu}_{k+1})) \\ & \leq L_2 \mathbb{E} \mathcal{W}_2(\tilde{\nu}_{k+1}, \tilde{\nu}^*(\mu_k)) \\ & \stackrel{(104)}{\leq} L_2 \sqrt{\frac{F(\tilde{\mu}_k, \tilde{\nu}^*(\mu_k)) - \mathbb{E} F(\tilde{\mu}_k, \tilde{\nu}_{k+1})}{2\xi}} \\ & \leq L_2 \sqrt{\frac{\sigma^{K_\nu} [F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)) - F(\tilde{\mu}_k, \tilde{\nu}_0)] + \eta^\nu \bar{\varepsilon}_\nu \cdot \frac{1-\sigma^{K_\nu}}{1-\sigma}}{2\xi}} \\ & \leq \frac{L_2}{L_0} \cdot \Delta, \end{aligned} \quad (143)$$

where (104) relates Wasserstein distance to the objective difference, the last but one inequality follows from Theorem 34, the last inequality holds by the definition of  $\Delta$  in (142) and the same choice of  $K_\mu$  in the proof for a desired  $\mathcal{J}_\mu(\tilde{\mu}_k, \tilde{\nu}_{k+1})$  ( $k \in [K_\mu]$ ). Therefore, the second term in (23) of Theorem 8 is proved.

To sum up, we conclude the proof of our main theorem and the sketch of proof ideas.  $\blacksquare$

We additionally remark that for our convergence rate to the approximate IFNE coincides with the finite-dimensional case regardless of the order of numbers of particles adopted. More clearly, we resort to writing

our result in terms of the  $\theta$ -IFNE for some  $\theta > 0$ . First, we can alternatively claim that there exists  $k \in [K_\mu]$  such that the following equivalent result holds,

$$\mathbb{E}\mathcal{J}_\mu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = \mathcal{O}\left(\left(\Delta + \sqrt{\bar{\varepsilon}_\mu} + 2\tilde{L}R + G_{\max}\right) \cdot \sqrt{\frac{2}{1/\eta^\mu - \tilde{L}/2}\left(2R(\Delta + \sqrt{\bar{\varepsilon}_\mu}) + \frac{M_G}{K_\mu}\right)}\right), \quad (144)$$

$$\mathbb{E}\mathcal{J}_\nu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = \mathcal{O}\left(\frac{L_2}{L_0} \cdot \Delta\right). \quad (145)$$

By fixing  $N_\mu$  and  $N_\nu$  for any  $\theta > 0$ , we observe that when we set  $K_\mu \geq K_\mu(\theta) = \mathcal{O}(\theta^{-2})$  and  $K_\nu \geq K_\nu(\theta) = \mathcal{O}(2 \log(\theta^{-1}))$  simultaneously, we are able to obtain a  $\theta$ -IFNE in expectation since

$$\mathbb{E}\mathcal{J}_\mu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) \leq \theta, \quad (146)$$

$$\mathbb{E}\mathcal{J}_\nu(\tilde{\mu}_k, \tilde{\nu}_{k+1}) \leq \theta, \quad (147)$$

which is exactly the order in [58], implying the PL-condition and smoothness in infinite-dimensional settings work similarly as in finite-dimensional problems.

On the other hand, when the order of  $N_\mu$  and  $N_\nu$  dominate, we can compute that when  $N_\mu \geq N_\mu(\theta) = \mathcal{O}(\theta^{-4})$  and  $N_\nu \geq N_\nu(\theta) = \mathcal{O}(\theta^{-4})$  for any  $\theta > 0$ , we obtain the  $\theta$ -IFNE in expectation. Such a result implies that we need more particles than timesteps to guarantee a given precision of the solution. This also implies that the statistical error induced by measure approximation in infinite-dimensional spaces is more prominent than the optimization error.

## Appendix F. Proof of Convergence to the Minimax Value of Two-Sided PL Games

### F.1. PL Condition on the Inner Maximization Functional

**Lemma 36** *For a two-sided PL-game depicted by Assumption 9, the functional  $G(\mu) = \max_{\nu \in \mathcal{M}} F(\mu, \nu)$  satisfies the  $\xi_1$ -PL condition.*

**Proof** Since  $F_\nu(\mu) = F(\mu, \nu)$  satisfies the  $\xi_1$ -PL condition for any given  $\nu \in \mathcal{M}(\mathcal{X}_\nu)$ , we have

$$\begin{aligned} 2\xi_1 \cdot \left( F(\mu, \nu^*(\mu)) - \min_{\tilde{\mu} \in \mathcal{M}(\mathcal{X}_\mu)} F(\tilde{\mu}, \nu^*(\mu)) \right) &\leq \langle \text{grad } F(\mu, \nu^*(\mu)), \text{grad } F(\mu, \nu^*(\mu)) \rangle_\mu \\ &= \langle \text{grad } G(\mu), \text{grad } G(\mu) \rangle_\mu, \end{aligned} \quad (148)$$

where the last equality follows from Lemma 33. On the other hand, we have by definition

$$\min_{\tilde{\mu} \in \mathcal{M}(\mathcal{X}_\mu)} F(\tilde{\mu}, \nu^*(\mu)) \leq \min_{\tilde{\mu} \in \mathcal{M}(\mathcal{X}_\mu)} \max_{\nu \in \mathcal{M}(\mathcal{X}_\nu)} F(\tilde{\mu}, \nu) = \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu). \quad (149)$$

By plugging (149) into (148), we arrive at

$$2\xi_1 \cdot \left( G(\mu) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) \right) \leq \langle \text{grad } G(\mu), \text{grad } G(\mu) \rangle_\mu, \quad (150)$$

which implies that  $G(\mu)$  satisfies  $\xi_1$ -PL condition. ■

### F.2. Proof of Theorem 10

**Proof** The proof is based on Lemma 34 while different from the proof of Theorem 8 by the fact that we are now able to bound the norm of the Riemannian gradient of  $G(\mu)$  from below. By the smoothness of  $G$  under

Lemma 33, we have

$$G(\tilde{\mu}_{k+1}) \leq G(\tilde{\mu}_k) + \langle \text{grad}_\mu F(\tilde{\mu}_k, \nu^*(\tilde{\mu}_k)), \text{Exp}_{\tilde{\mu}_k}^{-1}(\tilde{\mu}_{k+1}) \rangle_{\tilde{\mu}_k} + \frac{\tilde{L}}{2} \langle \text{Exp}_{\tilde{\mu}_k}^{-1}(\mu_{k+1}), \text{Exp}_{\tilde{\mu}_k}^{-1}(\mu_{k+1}) \rangle_{\tilde{\mu}_k} \quad (151)$$

$$\begin{aligned} & \stackrel{(132)}{=} G(\tilde{\mu}_k) - \eta^\mu \langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k} - \eta^\mu \langle \text{grad } G(\tilde{\mu}_k), \delta_k - \Delta_k \rangle_{\tilde{\mu}_k} \\ & + \frac{\tilde{L}}{2} \langle \text{grad } G(\tilde{\mu}_k) - \Delta_k + \delta_k, \text{grad } G(\tilde{\mu}_k) - \Delta_k + \delta_k \rangle_{\tilde{\mu}_k} \\ & \leq G(\tilde{\mu}_k) - \left( \eta^\mu - (\eta^\mu)^2 \cdot \frac{\tilde{L}}{2} \right) \cdot \langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k} \\ & + \left( \eta^\mu + \tilde{L}(\eta^\mu)^2 \right) \left| \langle \text{grad } G(\tilde{\mu}_k), \delta_k - \Delta_k \rangle_{\tilde{\mu}_k} \right| + \frac{(\eta^\mu)^2 \tilde{L}}{2} \langle \delta_k - \Delta_k, \delta_k - \Delta_k \rangle_{\tilde{\mu}_k}. \end{aligned} \quad (152)$$

Here  $\delta_k = \widehat{\text{grad}}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) - \text{grad}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1}) = -\text{div}[\tilde{p}_k \cdot (\nabla \tilde{f}_k^* - \nabla f_k^*)]$  and  $\Delta_k = \text{grad } G(\tilde{\mu}_k) - \text{grad}_\mu F(\tilde{\mu}_k, \tilde{\nu}_{k+1})$  are defined in (130). Since  $\langle \cdot, \cdot \rangle_{\tilde{\mu}_k}$  serves as the Riemannian metric of  $\mathcal{M}(\mathcal{X}_\mu)$  at  $\tilde{\mu}_k$ , then by the Cauchy-Schwartz inequality we obtain

$$\begin{aligned} 2 \left| \langle \text{grad } G(\tilde{\mu}_k), \delta_k - \Delta_k \rangle_{\tilde{\mu}_k} \right| & \leq \langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k} + \langle \delta_k - \Delta_k, \delta_k - \Delta_k \rangle_{\tilde{\mu}_k} \\ & \leq \langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k} + 2 \langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + 2 \langle \Delta_k, \Delta_k \rangle_{\tilde{\mu}_k}. \end{aligned} \quad (153)$$

Hence by plugging (153) into (151), we have

$$\begin{aligned} G(\tilde{\mu}_{k+1}) & \leq G(\tilde{\mu}_k) + \frac{\eta^\mu(2\eta^\mu\tilde{L} - 1)}{2} \cdot \langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k} \\ & + \eta^\mu \left( 1 + 2(\eta^\mu)^2\tilde{L} \right) \cdot \left( \langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + \langle \Delta_k, \Delta_k \rangle_{\tilde{\mu}_k} \right). \end{aligned} \quad (154)$$

Since  $\eta^\mu \leq 1/(2\tilde{L})$ , we require a lower bound on  $\langle \text{grad } G(\tilde{\mu}_k), \text{grad } G(\tilde{\mu}_k) \rangle_{\tilde{\mu}_k}$ . Following from the  $\xi_1$ -PL condition of  $G$  in Lemma 36, we can further obtain

$$\begin{aligned} G(\tilde{\mu}_{k+1}) & \leq G(\tilde{\mu}_k) + \xi_1 \eta^\mu (2\eta^\mu\tilde{L} - 1) \cdot \left( G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) \right) \\ & + \eta^\mu \left( 1 + 2(\eta^\mu)^2\tilde{L} \right) \cdot \left( \langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + \langle \Delta_k, \Delta_k \rangle_{\tilde{\mu}_k} \right), \end{aligned} \quad (155)$$

which can be rewritten as

$$\begin{aligned} G(\tilde{\mu}_{k+1}) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) & \leq [1 - \xi_1 \eta^\mu (1 - 2\eta^\mu\tilde{L})] \cdot \left( G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) \right) \\ & + \eta^\mu \left( 1 + 2(\eta^\mu)^2\tilde{L} \right) \cdot \left( \langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + \langle \Delta_k, \Delta_k \rangle_{\tilde{\mu}_k} \right). \end{aligned} \quad (156)$$

Now we invoke the upper bounds on  $\|\delta_k\|_{\tilde{\mu}_k}$  in (101) and  $\|\Delta_k\|_{\tilde{\mu}_k}$  in (136) to further develop a contraction for the value of  $G$  at each timestep,

$$\begin{aligned} \mathbb{E}G(\tilde{\mu}_{k+1}) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) & \leq [1 - \xi_1 \eta^\mu (1 - 2\eta^\mu\tilde{L})] \cdot \left( \mathbb{E}G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) \right) \\ & + \eta^\mu \left( 1 + 2\eta^\mu\tilde{L} \right) \cdot \left( \mathbb{E}\langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + \tilde{\Delta}^2 \right), \end{aligned} \quad (157)$$

where we define

$$\tilde{\Delta}^2 = L_0^2 \cdot \frac{\sigma^{K_\nu} \cdot M_H + \eta^\nu \bar{\varepsilon}_\nu \cdot \frac{1 - \sigma^{K_\nu}}{1 - \sigma}}{2\xi_2}, \quad (158)$$

with  $\xi_2$  as the parameter of PL condition for the inner-loop player,  $K_\nu$  as the number of timesteps for the inner-loop player defined in Algorithm 1, and the expectation is taken with respect to the randomness of initial particles. Also, by the definitions of  $\tilde{L}$ ,  $\xi_1$ , and  $\eta^\mu \in (0, 1/(4\tilde{L}))$ , we have  $1 - \xi_1\eta^\mu(1 - 2\eta^\mu\tilde{L}) < 1 - \xi_1\eta^\mu/2 < 1$ . For simplicity, let  $\tilde{\sigma} = 1 - \xi_1\eta^\mu/2$ , by multiplying  $\tilde{\sigma}^{-(k+1)}$  to (157) we have

$$\begin{aligned} \tilde{\sigma}^{-(k+1)} \cdot [\mathbb{E}G(\tilde{\mu}_{k+1}) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)] &\leq \tilde{\sigma}^{-k} \cdot [\mathbb{E}G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)] \\ &+ \tilde{\sigma}^{-(k+1)} \cdot \eta^\mu \cdot \left( \mathbb{E}\langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} + L_0^2 \cdot \frac{\eta^\nu \bar{\varepsilon}_\nu + \theta}{2\xi(1-\sigma)} \right), \end{aligned} \quad (159)$$

Note that  $\mathbb{E}\langle \delta_k, \delta_k \rangle_{\tilde{\mu}_k} = \bar{\varepsilon}_k$ , which is bounded in (101).. Then by summing up over the telescoping sequence  $\{\tilde{\sigma}^{-k} \cdot [\mathbb{E}G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)]\}_{k \geq 0}$  in (159), we obtain the following optimization error bound at iteration  $k$  for  $k \geq 1$ ,

$$\begin{aligned} \mathbb{E}G(\tilde{\mu}_k) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu) &\leq \tilde{\sigma}^k \cdot [G(\tilde{\mu}_0) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)] \\ &+ \sum_{m=0}^{k-1} \tilde{\sigma}^{k-(m+1)} \cdot \eta^\mu \cdot \left( \bar{\varepsilon}_m + L_0^2 \cdot \frac{\eta^\nu \bar{\varepsilon}_\nu + \theta}{2\xi(1-\sigma)} \right) \\ &\stackrel{(101)}{\leq} \tilde{\sigma}^k \cdot [G(\tilde{\mu}_0) - \min_{\mu \in \mathcal{M}(\mathcal{X}_\mu)} G(\mu)] \\ &+ \frac{1 - \tilde{\sigma}^k}{1 - \tilde{\sigma}} \cdot \eta^\mu \cdot \left( \bar{\varepsilon}_\mu + L_0^2 \cdot \frac{\eta^\nu \bar{\varepsilon}_\nu + \theta}{2\xi(1-\sigma)} \right), \end{aligned} \quad (160)$$

where

$$\begin{aligned} \bar{\varepsilon}_\mu &= \mathcal{O} \left( \tilde{K} \left( \frac{r_f^2}{N_\mu^{1/2}} + \frac{r_f^3}{w^{1/2}} + \frac{r_f^4}{w} \right) \right), \\ \tilde{\Delta}^2 &= L_0^2 \cdot \frac{\sigma^{K_\nu} \cdot M_H + \eta^\nu \bar{\varepsilon}_\nu \cdot \frac{1-\sigma^{K_\nu}}{1-\sigma}}{2\xi_2}. \end{aligned} \quad (161)$$

By the definition of  $G$ , we have  $\min_{\mu \in \mathcal{M}(\mathcal{X})} G(\mu) = F(\mu^*, \nu^*)$ . As a result, (160) gives the error bound in objective value  $F$  at iteration  $k$ . Hence we conclude the proof of Theorem 10.  $\blacksquare$

## Appendix G. Toy Experiments

In this section we report some results for a toy experiment with a Gaussian mixture model with 8 Gaussian distributions. For simplicity, we drop the regularizer terms from WGAN loss and consider a mixture of 8 generators and discriminators corresponding to the particles for parameters of the generator and the discriminator of WGAN. Both generators and discriminators are MLP with 3 layers. We also don't tune the learning rate and set it to be  $10^{-4}$ . We run the model for 20000 iterations which is small compared to the typical number of iterations used in practice to train a WGAN model. In our experiment we reused the code provided by [31] with some simple modification. We present some samples generated from trained generators in Figure 2. The blue dots are generated from real mixture models and the red ones are generated from generators. We observe that the distribution generated by our generator matches the groundtruth after a short training period.




Figure 2: Toy experiment results: blue dots represent samples from the gaussian mixture and the red dots represent the samples generated from generators.